

# Cancer in Tasmania

Incidence and Mortality 2002



Tasmania

DEPARTMENT of  
HEALTH and  
HUMAN SERVICES

---

# Cancer in Tasmania

## Incidence and Mortality

### 2002

---

#### Tasmanian Cancer Registry

University of Tasmania  
Menzies Research Institute  
17 Liverpool Street  
Hobart Tasmania Australia 7000

Postal address:

Private Bag 23  
Hobart Tasmania Australia 7001

Telephone (03) 6226 7757

Facsimile (03) 6226 7755

URL: [www.menzies.utas.edu.au/cancer/cancer.htm](http://www.menzies.utas.edu.au/cancer/cancer.htm)

Email: [tcr@menzies.utas.edu.au](mailto:tcr@menzies.utas.edu.au)

Editors: Newman L, Venn A, Albion T, Blizzard L, Fryer J

Publication date: May 2005



---

# Acknowledgements

---

The Tasmanian Cancer Registry is funded by the Tasmanian Department of Health and Human Services (DHHS), through its Population Health subdivision, and by the Menzies Research Institute. Additional funds are raised from the community and research funding bodies. The work of collecting and collating cancer registry data would not be possible without the continuing assistance and support of a number of people. These include staff members of:

- Private and public pathology laboratories
- The Registry of Births, Deaths and Marriages
- Medical Records Departments of all Tasmanian hospitals
- WP Holman Clinics in Launceston and Hobart
- Medical practitioners in specialist and general practices
- The Australian Bureau of Statistics
- Other State and Territory cancer registries

The assistance provided by the Registry's Advisory Committee is greatly appreciated. The administrative staff and volunteers of the Registry are commended for their commitment and efforts towards achieving the high level of accuracy and completeness of data upon which this report is based. We also thank other Menzies Research Institute staff for their assistance with statistical, computing, administrative, financial, media and editorial matters.

## Citation

The following citation is suggested in referring to this report:

Newman L, Venn A, Albion T, Blizzard L, Fryer J. Cancer in Tasmania: Incidence and Mortality 2002. Menzies Research Institute, Hobart, 2005.

---

# Tasmanian Cancer Registry Staff

---

Dr A Venn	Director
Ms L Newman	Manager
Mrs K Jackman	Administrative Officer
Mrs V Webb	Administrative Officer
Mrs P Whelan	Administration Assistant
Mrs J Clifford	Volunteer
Mrs M Magill	Volunteer
Dr L Blizzard	Biostatistician, Menzies Research Institute
Miss J Fryer	Biostatistician, Menzies Research Institute
Mr T Albion	IT Systems Manager, Menzies Research Institute
Mrs D Shugg	Honorary Research Associate

## Members of the Advisory Committee 2005

Mr K Churchill	Australian Bureau of Statistics, Tasmania
TBA	The Cancer Council Tasmania
Ms G Raw	Department of Health and Human Services, Tasmania
Professor P Stanton	School of Medicine, Department of Surgery, University of Tasmania, Tasmania
Dr R Taylor	Director of Public Health and Deputy Director of Population Health, Department of Health and Human Services, Tasmania
Dr R Young	School of Medicine, Department of Clinical Sciences, University of Tasmania, Tasmania



---

# Contents

---

<b>New reporting in ICD-10 codes .....</b>	<b>7</b>
<b>The Tasmanian Cancer Registry .....</b>	<b>8</b>
Introduction .....	8
Sources of data .....	8
Data handling, collection and coding practices .....	8
Data control and quality assurance .....	8
Publication of reports .....	9
<b>All Cancers .....</b>	<b>10</b>
Incidence and mortality of all cancers 2002 .....	10
Trends for all cancers 1980 – 2002 .....	11
<b>Common Cancers .....</b>	<b>12</b>
Common cancers diagnosed in males 2002 .....	12
Common cancers diagnosed in females 2002 .....	13
<b>Regional Distribution of Cancers .....</b>	<b>14</b>
<b>Female breast cancer 1997 - 2002 .....</b>	<b>15</b>
Tumour size .....	15
Lymph node involvement .....	15
<b>Cancer incidence and mortality table 2002.....</b>	<b>17</b>
Coding differences between ICD-9 & ICD-10 .....	54
<b>Appendices.....</b>	<b>55</b>
Appendix A: Cancer site codes and combinations.....	56
Appendix B: Statistical methods .....	58
Appendix C: Population data .....	59
Appendix D: Indices of data quality .....	60
Appendix E: Use of Tasmanian Cancer Registry Data .....	62
Appendix F: Incidence and mortality summary tables 2002.....	64



---

# New reporting in ICD-10 codes

---

This report provides a summary of cancer incidence and mortality statistics for Tasmania in 2002. It includes all cancers notified to the Tasmanian Cancer Registry that were first diagnosed in Tasmanian residents between 1 January 2002 and 31 December 2002.

The primary site of cancers diagnosed in Tasmania in 2002 was coded using the International Classification of Diseases for Oncology, ninth edition (ICD-9<sup>1</sup>), and the morphology was coded using ICD-O-2<sup>2</sup>.

In past annual reports, cancers were tabulated by ICD-9 codes, however in this report cancers have been tabulated according to ICD-10<sup>3</sup> codes (refer to Appendix A on page 56). ICD-9 and ICD-10 codes are comparable for most cancers. Where ICD-9 and ICD-10 codes are not comparable an explanation of these differences is provided on page 54.

This report presents data for invasive cancers only (behaviour = 3, site C000 – C809). In situ cancers and second primary cancers with the same three-digit topography code and related morphologies are not included in this report.

In past annual reports, age standardised rates were calculated using only the World Standard Population (1960). In this report, age standardised rates have also been calculated using the Australian Standard Population (2001) and the World Standard Population (2000) to enable Tasmania's cancer data to be compared with data published by the Australian Institute of Health and Welfare.

---

<sup>1</sup> International Classification of Diseases, 9<sup>th</sup> revision

<sup>2</sup> International Classification of Diseases for Oncology, 2<sup>nd</sup> edition

<sup>3</sup> International Classification of Diseases, 10<sup>th</sup> revision



# The Tasmanian Cancer Registry

## Introduction

The Tasmanian Cancer Registry was established in 1977 as a population-based registry covering the whole of Tasmania. The Registry was established to provide the State Government with accurate cancer incidence and mortality statistics and to monitor cancer trends. In July 1988 the operation of the Cancer Registry was moved from the Department of Health Services to the Menzies Research Institute. Cancer was proclaimed a notifiable disease in December 1992 and cancer registration since then has had a legislative basis.

Registry staff includes a Director, Manager, two Administrative Officers and an Administration Assistant. Volunteers also assist with the paper data handling. The Registry has access to a biostatistician and a computer consultant. An Advisory Committee assists the Registry. The Tasmanian Cancer Registry is a full member of the Australasian Association of Cancer Registries (AACR) and the International Association of Cancer Registries (IACR).

## Sources of data

All pathology laboratories in the State provide the Registry with copies of histopathological and cytology reports of cancer and cell marker reports. Cancer notification forms are supplied by the two radiation oncology clinics. Private and public hospitals notify diagnoses of cancer to the Registry upon discharge of patients or provide a computerised listing of cancer cases periodically. Death notifications of Tasmanian people are reviewed for mention of cancer as a cause of death. Since 1994 breast and cervical cancer screening programs have been undertaken in Tasmania and listings from these sources are available to check against Registry records. Interstate registries supply data to the Tasmanian Cancer Registry on Tasmanian residents who seek treatment interstate or who move interstate at some time after cancer diagnosis.

## Data handling, collection and coding practices

Paper copies of all information are retained. Paper records for persons deceased are archived two years after death. Data for cancers diagnosed between 1 January 2002 and 31 December 2002 were registered on database software provided by the

South Australian Cancer Registry. Only one tumour per ICD-9 site can be recorded. The Tasmanian Cancer Registry collects and stores all pathology reports of non-melanoma skin cancers (NMSC) but these are not routinely entered onto the database due to resource considerations. Deaths from NMSC are reported annually.

Information collected by the Registry includes demographic and clinical data for the cancer patient. Additional tumour data are collected for melanoma, lymphomas, unknown primaries, and breast and bladder cancers.

The primary site of cancers diagnosed in Tasmania in 2002 was coded to the International Classification of Diseases for Oncology, ninth edition (ICD-9<sup>4</sup>), and the morphology was coded to ICD-O-2<sup>5</sup>. This report presents data for invasive cancers only (behaviour = 3, site C000 – C809). In situ cancers and second primary cancers with the same three-digit topography code and related morphologies are not included in this report. Coding practices specific to the Tasmanian Cancer Registry are detailed in Appendix A. The Tasmanian Cancer Registry records multiple primary cancers in the same person, of which only some are counted for incidence purposes according to the rules of the International Agency for Research on Cancer (IARC) and IACR.

## Data control and quality assurance

The quality of information provided by the Registry depends on the quality of data received. The indices used to measure the quality of the 2002 data are provided in Appendix D. To help achieve high data quality and case ascertainment, data are obtained from multiple sources such as pathology laboratories, hospitals and the Registrar of Births, Deaths and Marriages. Most registered cases include data from both a pathology laboratory and a hospital service (inpatient or radiation oncology clinic). Where insufficient information is received to enable complete registration, active follow-up is undertaken by

---

<sup>4</sup> International Classification of Diseases, 9<sup>th</sup> revision

<sup>5</sup> International Classification of Diseases for Oncology, 2<sup>nd</sup> edition

contacting treating doctors, pathology laboratories and hospital medical records departments. The quality also depends on the accuracy of processing of data by the Registry. The existing system is able to detect a number of errors when data entry is performed. Data matching programs enable the identification and amendment of duplicate entries by identifying incorrect spellings, name changes and date of birth inconsistencies. In addition, the National Cancer Statistics Clearing House (NCSCH) collates all State and Territory data and checks for duplicate registrations across two or more States.

Notifications of cancer are received and dealt with in a timely manner at the Tasmanian Cancer Registry. On average, the Registry receives cancer notifications once or twice a week from pathology laboratories, quarterly and ad hoc from hospitals, and monthly from the Registrar of Births, Deaths and Marriages. The cases are registered within two months of notification and cases that are incomplete are followed up over the following 18 months.

## **Publication of reports**

The incidence and mortality data in this report are based on cancer registrations for 2002, and for 1978 - 2002 in trend analysis, completed by 30 September 2004. Despite intensive efforts to ensure the completeness of incidence data, the database is continually updated with previously unregistered cases and new information for registered cases. This improves the quality of data but future publications and responses to requests for data will reflect any subsequent revisions to the data and may not exactly correspond to the figures in this report.

# All Cancers


## Incidence and mortality of all cancers 2002

There were 2,534 new cases of cancer (excluding non-melanoma skin cancers) diagnosed among Tasmanian residents during 2002 (1,398 males and 1,136 females). The overall age standardised incidence was 374.4 per 100,000 for males and 285.8 per 100,000 for females<sup>6</sup>. The risk of developing any cancer by age 75 years was 1 in 3 for males and 1 in 4 for females. The risk estimate does not include the risk of developing non-melanoma skin cancer.

Cancer incidence generally increased with age (Figure 1). Male rates exceeded female rates for Tasmanians aged 60 years and over. Prostate cancer and lung cancer were responsible for the greater male cancer incidence at these ages. Breast cancer accounted for the slightly higher female rates among younger adults.

There were 1,128 (639 male and 489 female) cancer-related deaths among Tasmanian residents in 2002. Four deaths were due to non-melanoma skin cancer. The overall age standardised mortality rate was 158.7 per 100,000 for males and 102.0 per 100,000 for females. The person years life lost to age 75 years was 5,326 for males and 4,480 for females.

Figure 1: Age specific incidence and mortality for all cancers (excluding non-melanoma skin cancers) 2002




<sup>6</sup> Age standardised incidence was calculated using the World Standard Population (1960)

## Trends for all cancers 1980 – 2002

The age standardised incidence rates of all cancers (excluding non-melanoma skin cancers) increased by 33.0% for males and 33.2% for females during the 22-year period from 1980 to 2002 (comparing 2001-2002 with 1980-1981). Some of the largest increases were observed for prostate cancer in males and breast cancer in females. The increase in incidence during the 1990s coincided with the greater use of prostate-antigen testing (PSA) for prostate cancer in men and the introduction of mammography screening for breast cancer in women. While the number of new cancer cases diagnosed each year has risen over the period from 1980 to 2002, and the number of deaths from cancer has risen as well, age standardised mortality rates per 100,000 population have remained relatively stable. As can be seen from Figure 2, this difference has resulted in a widening between incidence and mortality rates.

**Figure 2: Trends in age standardised incidence and mortality of all cancers (excluding non-melanoma skin cancers) 1980 – 2002\***




\*Incidence and mortality for all cancers from 1980 to 2000 is reported in ICD-9 while data for 2001 and 2002 are reported in ICD-10.

# Common Cancers

## Common cancers diagnosed in males 2002

The most common cancer diagnosed in males in 2002 was prostate cancer, followed by lung cancer, colorectal cancer, melanoma skin cancer and bladder cancer. While the number of prostate cancer cases was much greater than any other cancer, prostate cancer numbers have declined overall since the peak in 1994 when 420 new prostate cancer cases were reported.


Figure 3: Common cancers diagnosed in males 2002



## Common cancer-related deaths in males 2002

The most common causes of cancer-related deaths in male Tasmanian residents in 2002 were lung cancer, prostate cancer, colorectal cancer, cancer of the pancreas and cancer of the oesophagus.


Figure 4: Common causes of cancer-related deaths in males 2002



## Common cancers diagnosed in females 2002

The most common cancer diagnosed in females in 2002 was breast cancer, followed by colorectal cancer, melanoma skin cancer, lung cancer and lymphoma.


Figure 5: Common cancers diagnosed in females 2002



## Common cancer-related deaths in females 2002

The most common causes of cancer-related deaths in female Tasmanian residents in 2002 were lung cancer, colorectal cancer, breast cancer, cancer of the pancreas and ovarian cancer.

Figure 6: Common causes of cancer-related deaths in females 2002



# Regional Distribution of Cancers

The regional distribution for each cancer site is shown as the number of cases and the percentage of all cases of each cancer site in each statistical division (Table 1). This information is based on recorded postcode of residence. On the basis of population numbers in each of the statistical divisions, the distribution of cancers would be expected to be 49% in the South, 28% in the North and 23% in the Mersey-Lyell division. Variation around that distribution can be expected due to chance occurrences and differences in the age distribution between the regional populations.



**Table 1: Regional distribution of cancer incidence for all sites with a minimum of 50 new cases 2002**

ICD-10	Site	Southern	Northern	Mersey-Lyell	Total*
		232,576 (49%)	133,644 (28%)	106,392 (23%)	
C18 - C21	Colorectal	166 (48%)	86 (25%)	91 (27%)	343
C50	Breast	168 (50%)	104 (31%)	61 (18%)	333
C61	Prostate	146 (47%)	93 (30%)	73 (23%)	312
C33, C34	Lung	135 (44%)	96 (32%)	73 (24%)	304
C43	Melanoma of skin	121 (54%)	59 (26%)	45 (20%)	225
C67	Bladder	65 (51%)	44 (35%)	18 (14%)	127
C81 - C85	All lymphomas	49 (50%)	29 (30%)	20 (20%)	98
C64	Kidney	35 (49%)	22 (31%)	15 (21%)	72
C25	Pancreas	33 (52%)	20 (31%)	11 (17%)	64
C91 - C95	All leukaemia	29 (48%)	17 (28%)	14 (23%)	60
C01-C14,C30-C32	Head and neck	29 (54%)	17 (31%)	8 (15%)	54
C16	Stomach	20 (38%)	16 (30%)	17 (32%)	53
C15	Oesophagus	19 (46%)	10 (24%)	12 (29%)	41
C71	Brain	22 (59%)	11 (30%)	4 (11%)	37
C56	Ovary	16 (57%)	6 (21%)	6 (21%)	28
<b>Total new cases</b>		<b>1,053 (49%)</b>	<b>630 (29%)</b>	<b>468 (22%)</b>	<b>2,151</b>

\*Source: Australian Bureau of Statistics, Regional Population Growth, Australia and New Zealand, 2002-03, (cat. no. 3218), pp 83-84.


# Female breast cancer 1997 - 2002

The Tasmanian Cancer Registry first recorded breast cancer tumour size and lymph node involvement in 1997 when funding was provided to all Australian cancer registries for this purpose.

## Tumour size

In 2002 97% of the 333 primary breast cancer cases (female) were histologically examined<sup>7</sup>. Information about tumour size was available for 304 (91.0%) of these cases. Of these tumours, 57 (18.8%) were less than 10mm in diameter, 120 (39.5%) were between 10 and 19mm, 112 (36.8%) were between 20 and 49mm, and 15 (4.9%) were greater than 50mm in diameter. Figure 7 compares categories of tumour size from 1997 to 2002.


Figure 7: Breast cancer - Size of histologically confirmed tumours 1997 - 2002



## Lymph node involvement

Of the 333 primary breast cancer cases (female), lymph nodes were investigated in 220 (66.1%) of cases. Where nodal status was examined, 119 (54.1%) cases were classified as lymph node negative, 67 (30.5%) cases involved 1 to 3 lymph nodes, 14 (6.4%) cases involved 4 to 6 lymph nodes, and 20 (9.1%) cases involved 7 or more lymph nodes. Figure 8 compares categories of lymph node involvement from 1997 to 2002.

Figure 8: Breast cancer - Lymph node involvement 1997 - 2002



<sup>7</sup> Refer to Indices of Data Quality on page 60.





---

# Cancer incidence and mortality table 2002

---

- Numbers of new cases and deaths
- Age-specific incidence and mortality rates per 100,000
- Crude incidence and mortality rates per 100,000 (Crude Rates)
- Cumulative incidence and mortality rates (Cumul Rates)
- Age standardised incidence and mortality rates using the Australian standard population (2001) and the world standard population (1960)



# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C00. Lip</b>																									
Cases		M	0	0	0	0	1	0	0	2	1	2	2	5	1	1	0	7	0	4	26				
		F	0	0	0	0	0	0	0	0	2	0	0	0	2	1	0	2	4	1	12				
		P	0	0	0	0	1	0	0	2	3	2	2	5	3	2	0	9	4	5	38				
Incidence per 100,000		M	0	0	0	0	6.9	0	0	12.5	5.5	11.8	12.3	35.6	8.8	10.9	0	112.3	0	175.0	11.2	0.5	7.1	11.5	
		F	0	0	0	0	0	0	0	0	10.8	0	0	0	17.8	10.6	0	25.9	69.6	19.5	5.0	0.2	2.4	4.1	
		P	0	0	0	0	3.5	0	0	6.1	8.2	5.9	6.2	17.9	13.3	10.7	0	64.5	43.3	67.5	8.0	0.4	4.6	7.4	
Deaths		M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
		F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
		P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mortality per 100,000		M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
		F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
		P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
<b>C01, C02. Tongue</b>																									
Cases		M	0	0	0	0	0	0	0	0	0	0	4	2	2	2	0	2	0	12					
		F	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1					
		P	0	0	0	0	0	0	0	1	0	0	4	2	2	2	0	2	0	13					
Incidence per 100,000		M	0	0	0	0	0	0	0	0	0	0	28.5	17.5	21.8	24.8	0	57.3	0	5.2	0.5	3.3	4.8		
		F	0	0	0	0	0	0	0	5.4	0	0	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4		
		P	0	0	0	0	0	0	0	2.7	0	0	14.3	8.8	10.7	11.8	0	21.7	0	2.8	0.2	1.8	2.5		
Deaths		M	0	0	0	0	0	0	1	0	0	0	1	0	1	2	0	0	1	6					
		F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
		P	0	0	0	0	0	0	1	0	0	0	1	0	1	2	0	0	1	6					
Mortality per 100,000		M	0	0	0	0	0	0	6.2	0	0	0	7.1	0	10.9	24.8	0	0	43.7	2.6	0.2	1.7	2.6		
		F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
		P	0	0	0	0	0	0	3.0	0	0	0	3.6	0	5.4	11.8	0	0	13.5	1.3	0.1	0.8	1.2		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C03. Gum</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1			
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	0	0	0	0	0.4	0.1	0.3	0.4
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	5.4	0	0	0	0	0	0.2	0.0	0.2	0.2
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
<b>C04. Floor of mouth</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	3			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	3			
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	6.2	0	8.8	0	0	16.0	0	0	1.3	0.1	0.8	1.2	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	3.1	0	4.4	0	0	7.2	0	0	0.6	0.0	0.4	0.6	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0	0	0.4	0.1	0.2	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	0	0	0	0.2	0.0	0.1	0.2	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C05, C06. Other mouth</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	3				
	F	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	2				
	P	0	0	0	0	0	0	1	0	0	0	1	0	1	0	1	0	1	0	5				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	6.2	0	8.8	0	12.4	0	0	0	1.3	0.1	0.9	1.2	
	F	0	0	0	0	0	0	6.1	0	0	0	0	0	0	0	0	0	17.4	0	0.8	0.0	0.5	0.8	
	P	0	0	0	0	0	0	3.1	0	0	0	3.1	0	4.4	0	5.9	0	10.8	0	1.1	0.1	0.7	1.0	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	2				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	3				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	10.9	0	16.0	0	0	0.9	0.1	0.5	0.8	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	0	0	0	0.4	0.1	0.3	0.4	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	10.7	0	7.2	0	0	0.6	0.1	0.4	0.6	
<b>C01 - C06. Oral Cavity</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	2	4	4	2	3	1	2	0	18				
	F	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	1	0	4				
	P	0	0	0	0	0	0	1	0	1	0	2	4	4	3	3	1	3	0	22				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	12.3	28.5	35.1	21.8	37.2	16.0	57.3	0	7.7	0.7	5.0	7.2	
	F	0	0	0	0	0	0	6.1	0	5.4	0	0	0	0	10.6	0	0	17.4	0	1.7	0.1	1.1	1.5	
	P	0	0	0	0	0	0	3.1	0	2.7	0	6.2	14.3	17.7	16.1	17.8	7.2	32.5	0	4.7	0.4	3.0	4.2	
Deaths																								
	M	0	0	0	0	0	0	0	1	0	0	0	1	0	2	3	1	0	1	9				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1				
	P	0	0	0	0	0	0	0	1	0	0	0	1	0	3	3	1	0	1	10				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	6.2	0	0	0	7.1	0	21.8	37.2	16.0	0	43.7	3.9	0.4	2.4	3.9	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	0	0	0	0.4	0.1	0.3	0.4	
	P	0	0	0	0	0	0	0	3.0	0	0	0	3.6	0	16.1	17.8	7.2	0	13.5	2.1	0.2	1.3	1.9	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C07, C08. Salivary glands</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	2				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0	0	0.4	0.1	0.2	0.4	
	F	0	0	0	0	0	0	0	0	0	5.8	0	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4	
	P	0	0	0	0	0	0	0	0	0	2.9	0	0	0	0	5.9	0	0	0	0.4	0.0	0.3	0.4	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	43.7	0.4	0.0	0.2	0.6	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.5	0.2	0.0	0.1	0.2	
<b>C09, C10. Oropharynx</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	1	0	0	4				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	1	0	0	4				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	5.5	0	6.2	0	8.8	0	0	16.0	0	0	1.7	0.1	1.2	1.6	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	2.7	0	3.1	0	4.4	0	0	7.2	0	0	0.8	0.1	0.6	0.8	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	2				
	P	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	2				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	0	0	11.3	0	0	0	0.8	0.1	0.5	0.8	
	P	0	0	0	0	0	0	0	0	0	0	3.1	0	0	0	5.9	0	0	0	0.4	0.0	0.3	0.4	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C11. Nasopharynx</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1			
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	19.5		0.4	0.0	0.1	0.3
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.5		0.2	0.0	0.1	0.2

## C12,C13. Hypopharynx

Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	2				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	2				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	28.6	0		0.9	0.1	0.4	0.9
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	0	10.8	0		0.4	0.0	0.2	0.4
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	2				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0	0		0.4	0.1	0.2	0.4
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	17.4	0		0.4	0.0	0.1	0.3
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	0	10.8	0		0.4	0.0	0.2	0.4



# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C09 - C13. Pharynx</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	1	0	1	0	1	0	1	1	1	0	6					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	1	0	1	0	1	0	1	1	1	0	6					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	5.5	0	6.2	0	8.8	0	12.4	16.0	28.6	0	2.6	0.2	1.5	2.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	2.7	0	3.1	0	4.4	0	5.9	7.2	10.8	0	1.3	0.1	0.7	1.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1					
	F	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	1	4					
	P	0	0	0	0	0	0	0	0	0	0	1	0	0	0	2	0	1	1	5					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0	0	0.4	0.1	0.2	0.4		
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	0	0	11.3	0	17.4	19.5	1.7	0.1	0.7	1.3		
	P	0	0	0	0	0	0	0	0	0	0	3.1	0	0	0	11.8	0	10.8	13.5	1.1	0.1	0.5	1.0		
<b>C14. Other oral</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	16.0	0	0	0.4	0.0	0.2	0.4		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.2	0	0	0.2	0.0	0.1	0.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C15. Oesophagus</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	2	3	4	3	9	6	2	0	29					
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	5	3	2	12					
	P	0	0	0	0	0	0	0	0	0	0	2	3	5	3	10	11	5	2	41					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	12.3	21.4	35.1	32.8	111.6	96.2	57.3	0	12.4	1.1	7.3	11.8		
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	0	11.3	64.9	52.2	39.1	5.0	0.1	1.7	3.9		
	P	0	0	0	0	0	0	0	0	0	0	6.2	10.8	22.1	16.1	59.2	78.9	54.1	27.0	8.7	0.6	4.5	7.8		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	3	2	6	4	6	6	3	1	31					
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	0	0	3	7					
	P	0	0	0	0	0	0	0	0	0	0	3	3	7	5	7	6	3	4	38					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	18.5	14.2	52.6	43.7	74.4	96.2	85.9	43.7	13.3	1.0	8.0	12.8		
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	10.6	11.3	0	0	58.6	2.9	0.2	1.5	2.3		
	P	0	0	0	0	0	0	0	0	0	0	9.2	10.8	30.9	26.9	41.5	43.0	32.5	54.0	8.0	0.6	4.6	7.2		
<b>C16. Stomach</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	1	2	1	5	3	4	12	2	2	32					
	F	0	0	0	0	0	0	1	1	0	2	2	0	2	1	1	5	5	1	21					
	P	0	0	0	0	0	0	1	1	0	3	4	1	7	4	5	17	7	3	53					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	12.3	7.1	43.9	32.8	49.6	192.5	57.3	87.5	13.7	0.8	7.6	13.6		
	F	0	0	0	0	0	0	6.1	5.9	0	11.6	12.3	0	17.8	10.6	11.3	64.9	87.0	19.5	8.8	0.4	4.5	7.5		
	P	0	0	0	0	0	0	3.1	3.0	0	8.8	12.3	3.6	30.9	21.5	29.6	121.9	75.8	40.5	11.2	0.6	5.9	10.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	1	1	1	3	1	2	6	2	5	22					
	F	0	0	0	0	0	0	1	0	1	0	0	3	1	3	2	4	1	1	17					
	P	0	0	0	0	0	0	1	0	1	1	1	4	4	4	4	10	3	6	39					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	7.1	26.3	10.9	24.8	96.2	57.3	218.7	9.4	0.4	5.2	10.0		
	F	0	0	0	0	0	0	6.1	0	5.4	0	0	21.6	8.9	31.8	22.7	51.9	17.4	19.5	7.1	0.5	4.0	6.2		
	P	0	0	0	0	0	0	3.1	0	2.7	2.9	3.1	14.3	17.7	21.5	23.7	71.7	32.5	81.0	8.3	0.4	4.4	7.5		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W (1960)	AUS (2001)	
<b>C17. Small intestine</b>																									
Cases																									
	M	0	1	0	0	0	0	0	0	0	1	1	0	0	1	1	0	1	2	8					
	F	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	2	0	0	5					
	P	0	1	0	0	1	0	0	0	0	1	1	1	0	1	2	2	1	2	13					
Incidence per 100,000																									
	M	0	5.9	0	0	0	0	0	0	0	5.9	6.2	0	0	10.9	12.4	0	28.6	87.5	3.4	0.2	2.4	3.7		
	F	0	0	0	0	7.1	0	0	0	0	0	0	7.2	0	0	11.3	25.9	0	0	2.1	0.1	1.3	1.9		
	P	0	3.0	0	0	3.5	0	0	0	0	2.9	3.1	3.6	0	5.4	11.8	14.3	10.8	27.0	2.8	0.2	1.8	2.6		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2	4					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	2	5					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	28.6	87.5	1.7	0.1	0.8	2.1		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0.4	0.0	0.1	0.3		
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	7.2	10.8	27.0	1.1	0.0	0.4	0.9		
<b>C18. Colon</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	1	1	5	9	15	16	26	20	15	10	118					
	F	0	0	0	0	0	1	0	0	2	3	6	5	6	20	18	21	9	18	109					
	P	0	0	0	0	0	1	0	0	3	4	11	14	21	36	44	41	24	28	227					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	5.5	5.9	30.9	64.1	131.6	174.7	322.5	320.8	429.4	437.4	50.7	3.7	29.3	50.4		
	F	0	0	0	0	0	7.2	0	0	10.8	17.4	36.9	36.1	53.4	211.7	204.2	272.4	156.7	351.7	45.5	2.9	23.4	38.1		
	P	0	0	0	0	0	3.6	0	0	8.2	11.7	33.9	50.2	92.8	193.5	260.7	294.0	259.8	378.2	48.0	3.3	26.0	43.3		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	1	1	2	8	4	8	11	3	8	46					
	F	0	0	0	0	0	0	1	2	2	0	1	3	4	2	3	9	7	15	49					
	P	0	0	0	0	0	0	1	2	2	1	2	5	12	6	11	20	10	23	95					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	14.2	70.2	43.7	99.2	176.4	85.9	350.0	19.7	1.2	11.3	20.3		
	F	0	0	0	0	0	0	6.1	11.8	10.8	0	6.2	21.6	35.6	21.2	34.0	116.7	121.9	293.1	20.4	0.7	8.9	16.3		
	P	0	0	0	0	0	0	3.1	6.1	5.5	2.9	6.2	17.9	53.0	32.2	65.2	143.4	108.3	310.6	20.1	1.0	10.0	18.1		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C19 - C21. Rectum</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	2	6	5	8	15	9	13	7	8	3	76				
	F	0	0	0	0	0	0	0	0	0	4	1	2	4	3	6	8	10	2	40				
	P	0	0	0	0	0	0	0	0	2	10	6	10	19	12	19	15	18	5	116				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	11.1	35.5	30.9	57.0	131.6	98.3	161.2	112.3	229.0	131.2	32.6	2.6	21.0	31.4	
	F	0	0	0	0	0	0	0	0	0	23.2	6.2	14.4	35.6	31.8	68.1	103.8	174.1	39.1	16.7	0.9	8.1	13.9	
	P	0	0	0	0	0	0	0	0	5.5	29.3	18.5	35.8	83.9	64.5	112.6	107.6	194.9	67.5	24.5	1.8	14.4	22.2	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	3	0	0	1	6	5	6	4	6	31				
	F	0	0	0	0	0	0	0	0	0	0	1	3	4	1	7	7	7	7	37				
	P	0	0	0	0	0	0	0	0	0	3	1	3	5	7	12	13	11	13	68				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	17.7	0	0	8.8	65.5	62.0	96.2	114.5	262.5	13.3	0.8	7.5	14.1	
	F	0	0	0	0	0	0	0	0	0	0	6.2	21.6	35.6	10.6	79.4	90.8	121.9	136.8	15.4	0.8	6.7	12.4	
	P	0	0	0	0	0	0	0	0	0	8.8	3.1	10.8	22.1	37.6	71.1	93.2	119.1	175.6	14.4	0.8	7.0	12.9	
<b>C18 - C21. Colorectal</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	3	7	10	17	30	25	39	27	23	13	194				
	F	0	0	0	0	0	1	0	0	2	7	7	7	10	23	24	29	19	20	149				
	P	0	0	0	0	0	1	0	0	5	14	17	24	40	48	63	56	42	33	343				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	16.6	41.4	61.7	121.1	263.2	272.9	483.7	433.0	658.5	568.7	83.3	6.3	50.3	81.8	
	F	0	0	0	0	0	7.2	0	0	10.8	40.6	43.1	50.5	89.0	243.5	272.2	376.2	330.8	390.8	62.2	3.8	31.5	52.0	
	P	0	0	0	0	0	3.6	0	0	13.6	41.0	52.4	86.0	176.7	258.0	373.2	401.6	454.7	445.7	72.6	5.0	40.4	65.5	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	4	1	2	9	10	13	17	7	14	77				
	F	0	0	0	0	0	0	1	2	2	0	2	6	8	3	10	16	14	22	86				
	P	0	0	0	0	0	0	1	2	2	4	3	8	17	13	23	33	21	36	163				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	23.7	6.2	14.2	78.9	109.2	161.2	272.7	200.4	612.4	33.1	2.0	18.7	34.4	
	F	0	0	0	0	0	0	6.1	11.8	10.8	0	12.3	43.3	71.2	31.8	113.4	207.5	243.7	429.9	35.9	1.5	15.6	28.7	
	P	0	0	0	0	0	0	3.1	6.1	5.5	11.7	9.2	28.7	75.1	69.9	136.3	236.7	227.3	486.2	34.5	1.7	17.0	31.0	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W (1960)	AUS (2001)	
<b>C22. Liver</b>																									
Cases																									
	M	0	0	0	0	0	0	0	1	0	1	3	0	2	0	2	0	0	0	9					
	F	0	0	0	0	0	0	0	0	0	0	0	0	2	2	1	1	1	0	7					
	P	0	0	0	0	0	0	0	1	0	1	3	0	4	2	3	1	1	0	16					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	6.2	0	5.9	18.5	0	17.5	0	24.8	0	0	0	3.9	0.4	2.9	3.7		
	F	0	0	0	0	0	0	0	0	0	0	0	0	17.8	21.2	11.3	13.0	17.4	0	2.9	0.3	1.8	2.5		
	P	0	0	0	0	0	0	0	3.0	0	2.9	9.2	0	17.7	10.7	17.8	7.2	10.8	0	3.4	0.3	2.3	3.1		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	2	0	2	1	3	0	0	0	8					
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	1	0	4					
	P	0	0	0	0	0	0	0	0	0	0	2	0	3	2	3	1	1	0	12					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	12.3	0	17.5	10.9	37.2	0	0	0	3.4	0.4	2.4	3.2		
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	10.6	0	13.0	17.4	0	1.7	0.1	0.9	1.4		
	P	0	0	0	0	0	0	0	0	0	0	6.2	0	13.3	10.7	17.8	7.2	10.8	0	2.5	0.2	1.6	2.3		
<b>C23, C24. Gallbladder</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	1	1	0	1	2	2	2	1	2	12					
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	2	0	1	1	4	10					
	P	0	0	0	0	0	0	0	0	0	1	1	1	2	4	2	3	2	6	22					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	0	8.8	21.8	24.8	32.1	28.6	87.5	5.2	0.3	3.1	5.3		
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	21.2	0	13.0	17.4	78.2	4.2	0.2	1.9	3.2		
	P	0	0	0	0	0	0	0	0	0	2.9	3.1	3.6	8.8	21.5	11.8	21.5	21.7	81.0	4.7	0.3	2.4	4.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	2	0	0	2	1	3	0	2	10					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	2	3	1	7					
	P	0	0	0	0	0	0	0	0	0	0	2	0	0	3	1	5	3	3	17					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	12.3	0	0	21.8	12.4	48.1	0	87.5	4.3	0.2	2.4	4.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	25.9	52.2	19.5	2.9	0.1	0.9	2.2		
	P	0	0	0	0	0	0	0	0	0	0	6.2	0	0	16.1	5.9	35.9	32.5	40.5	3.6	0.1	1.6	3.2		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C25. Pancreas</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	1	1	1	3	5	6	4	5	1	27					
	F	0	0	0	0	0	0	0	0	1	3	0	1	3	5	9	3	7	5	37					
	P	0	0	0	0	0	0	0	0	1	4	1	2	6	10	15	7	12	6	64					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	7.1	26.3	54.6	74.4	64.2	143.1	43.7	11.6	0.9	6.7	11.4		
	F	0	0	0	0	0	0	0	0	5.4	17.4	0	7.2	26.7	52.9	102.1	38.9	121.9	97.7	15.4	1.1	7.8	12.8		
	P	0	0	0	0	0	0	0	0	2.7	11.7	3.1	7.2	26.5	53.7	88.9	50.2	129.9	81.0	13.5	1.0	7.3	12.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	2	0	2	5	4	4	7	4	4	1	33					
	F	0	0	0	0	0	0	0	0	0	1	0	0	1	3	5	3	7	5	25					
	P	0	0	0	0	0	0	0	0	2	1	2	5	5	7	12	7	11	6	58					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	11.1	0	12.3	35.6	35.1	43.7	86.8	64.2	114.5	43.7	14.2	1.1	8.6	13.7		
	F	0	0	0	0	0	0	0	0	0	5.8	0	0	8.9	31.8	56.7	38.9	121.9	97.7	10.4	0.5	4.3	8.2		
	P	0	0	0	0	0	0	0	0	5.5	2.9	6.2	17.9	22.1	37.6	71.1	50.2	119.1	81.0	12.3	0.8	6.5	11.0		
<b>C30, C31. Nasal cavities</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	1	0	1	1	4	0	0	1	8					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	1	0	1	1	4	0	0	1	8					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	6.2	0	8.8	10.9	49.6	0	0	43.7	3.4	0.4	2.2	3.4		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0		
	P	0	0	0	0	0	0	0	0	0	0	3.1	0	4.4	5.4	23.7	0	0	13.5	1.7	0.2	1.0	1.5		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0	3					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0	3					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	14.2	0	10.9	0	0	0	0	0	1.3	0.1	0.9	1.1		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0		
	P	0	0	0	0	0	0	0	0	0	0	7.2	0	5.4	0	0	0	0	0	0.6	0.1	0.4	0.6		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C32. Larynx</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	2	4	3	1	1	1	0	12					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	1	0	3					
	P	0	0	0	0	0	0	0	0	0	0	0	2	4	5	1	1	2	0	15					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	14.2	35.1	32.8	12.4	16.0	28.6	0	5.2	0.5	3.5	4.7			
	F	0	0	0	0	0	0	0	0	0	0	0	0	21.2	0	0	17.4	0	1.3	0.1	0.7	1.0			
	P	0	0	0	0	0	0	0	0	0	0	7.2	17.7	26.9	5.9	7.2	21.7	0	3.2	0.3	2.1	2.8			
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	10.9	0	0	28.6	0	0.9	0.1	0.5	0.9		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	5.4	0	0	10.8	0	0.4	0.0	0.2	0.4		
<b>C33, C34. Lung</b>																									
Cases																									
	M	0	0	0	0	0	0	1	0	4	3	5	16	18	35	49	36	20	12	199					
	F	0	0	0	0	0	1	0	1	2	3	4	10	14	14	20	16	15	5	105					
	P	0	0	0	0	0	1	1	1	6	6	9	26	32	49	69	52	35	17	304					
Incidence per 100,000																									
	M	0	0	0	0	0	0	6.4	0	22.1	17.7	30.9	114.0	157.9	382.1	607.7	577.4	572.6	524.9	85.4	6.7	50.1	83.9		
	F	0	0	0	0	0	7.2	0	5.9	10.8	17.4	24.6	72.1	124.6	148.2	226.9	207.5	261.1	97.7	43.8	3.2	24.6	37.7		
	P	0	0	0	0	0	3.6	3.1	3.0	16.4	17.6	27.7	93.2	141.4	263.4	408.8	372.9	378.9	229.6	64.3	4.9	36.3	58.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	1	1	5	11	12	24	37	34	17	10	152					
	F	0	0	0	0	0	0	0	1	2	3	7	13	14	12	20	17	16	7	112					
	P	0	0	0	0	0	0	0	1	3	4	12	24	26	36	57	51	33	17	264					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	5.5	5.9	30.9	78.4	105.3	262.0	458.9	545.3	486.7	437.4	65.3	4.7	36.7	64.6		
	F	0	0	0	0	0	0	0	5.9	10.8	17.4	43.1	93.8	124.6	127.0	226.9	220.5	278.6	136.8	46.7	3.2	25.6	40.0		
	P	0	0	0	0	0	0	0	3.0	8.2	11.7	37.0	86.0	114.9	193.5	337.7	365.7	357.3	229.6	55.9	4.0	30.4	50.4		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C37, C38. Thymus etc</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	2				
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	7.1	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0	0.4	0.0	0.1	0.3	
	P	0	0	0	0	0	0	0	0	0	0	0	3.6	0	0	0	7.2	0	0	0	0.4	0.0	0.2	0.4	
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1				
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	7.1	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	3.6	0	0	0	0	0	0	0	0.2	0.0	0.1	0.2	
<b>C40, C41. Bone</b>																									
Cases																									
	M	1	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	3				
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1				
	P	1	0	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	4				
Incidence per 100,000																									
	M	6.4	0	0	0	0	0	0	0	0	0	0	7.1	0	0	12.4	0	0	0	0	1.3	0.1	1.3	1.2	
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	0	0	0	0	0	0	0.4	0.0	0.4	0.4	
	P	3.3	0	0	0	0	0	0	0	0	0	0	3.6	4.4	0	5.9	0	0	0	0	0.8	0.1	0.8	0.8	
Deaths																									
	M	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1				
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	5.5	0	0	0	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	2.7	0	0	0	0	0	0	0	0	0	0	0.2	0.0	0.2	0.2	



# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C43. Melanoma of skin</b>																								
Cases																								
	M	0	0	0	1	1	6	5	4	7	5	17	16	10	11	17	11	4	2	117				
	F	0	0	0	1	1	6	2	9	6	16	10	8	5	12	16	4	8	4	108				
	P	0	0	0	2	2	12	7	13	13	21	27	24	15	23	33	15	12	6	225				
Incidence per 100,000																								
	M	0	0	0	5.8	6.9	44.4	32.2	25.0	38.7	29.6	104.9	114.0	87.7	120.1	210.8	176.4	114.5	87.5	50.2	4.1	36.1	49.1	
	F	0	0	0	6.0	7.1	42.9	12.3	53.0	32.3	92.7	61.6	57.7	44.5	127.0	181.5	51.9	139.3	78.2	45.1	3.6	32.2	42.2	
	P	0	0	0	5.9	7.0	43.7	22.0	39.4	35.4	61.5	83.2	86.0	66.3	123.6	195.5	107.6	129.9	81.0	47.6	3.8	34.1	45.6	
Deaths																								
	M	0	0	0	0	1	0	0	0	1	1	1	2	2	0	6	5	4	1	24				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	3	5				
	P	0	0	0	0	1	0	0	0	1	1	1	2	2	0	6	6	5	4	29				
Mortality per 100,000																								
	M	0	0	0	0	6.9	0	0	0	5.5	5.9	6.2	14.2	17.5	0	74.4	80.2	114.5	43.7	10.3	0.7	5.9	10.3	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	17.4	58.6	2.1	0.0	0.5	1.4	
	P	0	0	0	0	3.5	0	0	0	2.7	2.9	3.1	7.2	8.8	0	35.5	43.0	54.1	54.0	6.1	0.3	3.1	5.6	
<b>C44. Skin</b>																								
Cases																								
	M	0	0	0	0	0	1	0	0	0	0	0	2	0	0	2	2	3	0	10				
	F	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	2	0	4				
	P	0	0	0	0	0	1	0	0	0	0	0	3	0	1	2	2	5	0	14				
Incidence per 100,000																								
	M	0	0	0	0	0	7.4	0	0	0	0	0	14.2	0	0	24.8	32.1	85.9	0	4.3	0.2	2.4	4.4	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	0	10.6	0	0	34.8	0	1.7	0.1	0.8	1.3	
	P	0	0	0	0	0	3.6	0	0	0	0	0	10.8	0	5.4	11.8	14.3	54.1	0	3.0	0.2	1.5	2.7	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	1	0	3				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	1	1	0	4				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	17.5	0	0	0	28.6	0	1.3	0.1	0.8	1.2	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0.4	0.0	0.1	0.3	
	P	0	0	0	0	0	0	0	0	0	0	0	0	8.8	0	0	7.2	10.8	0	0.8	0.0	0.5	0.8	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C45. Mesothelioma</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	1	3	0	3	2	3	0	0	12				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	1	3	0	4	2	3	0	0	13				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	6.2	21.4	0	32.8	24.8	48.1	0	0	5.2	0.4	3.1	4.8	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	0	0	0	0.4	0.1	0.3	0.4	
	P	0	0	0	0	0	0	0	0	0	0	3.1	10.8	0	21.5	11.8	21.5	0	0	2.8	0.2	1.7	2.5	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	1	4	0	3	2	3	1	0	14				
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	3				
	P	0	0	0	0	0	0	0	0	0	0	1	4	1	4	2	4	1	0	17				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	6.2	28.5	0	32.8	24.8	48.1	28.6	0	6.0	0.5	3.6	5.6	
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	10.6	0	13.0	0	0	1.3	0.1	0.8	1.1	
	P	0	0	0	0	0	0	0	0	0	0	3.1	14.3	4.4	21.5	11.8	28.7	10.8	0	3.6	0.3	2.1	3.2	
<b>C46. Kaposi sarcoma</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C48. Peritoneum</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
<b>C47, C49. Connective tissue</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	2	1	1	0	1	1	4	1	0	0	11				
	F	0	0	1	0	0	0	0	1	0	0	3	1	0	1	3	3	0	1	14				
	P	0	0	1	0	0	0	0	1	2	1	4	1	1	2	7	4	0	1	25				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	11.1	5.9	6.2	0	8.8	10.9	49.6	16.0	0	0	4.7	0.5	3.2	4.5	
	F	0	0	6.0	0	0	0	0	5.9	0	0	18.5	7.2	0	10.6	34.0	38.9	0	19.5	5.8	0.4	3.6	5.3	
	P	0	0	2.9	0	0	0	0	3.0	5.5	2.9	12.3	3.6	4.4	10.7	41.5	28.7	0	13.5	5.3	0.4	3.4	5.0	
Deaths																								
	M	0	0	1	1	0	1	0	1	0	1	1	0	1	0	0	0	0	0	7				
	F	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	1	1	0	4				
	P	0	0	1	2	0	1	0	1	0	1	1	0	2	0	0	1	1	0	11				
Mortality per 100,000																								
	M	0	0	5.7	5.8	0	7.4	0	6.2	0	5.9	6.2	0	8.8	0	0	0	0	0	3.0	0.2	3.0	3.0	
	F	0	0	0	6.0	0	0	0	0	0	0	0	0	8.9	0	0	13.0	17.4	0	1.7	0.1	1.1	1.4	
	P	0	0	2.9	5.9	0	3.6	0	3.0	0	2.9	3.1	0	8.8	0	0	7.2	10.8	0	2.3	0.2	2.1	2.3	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C50. Breast</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	2	11	12	26	43	40	44	36	38	29	18	21	13	333					
	P	0	0	0	0	0	2	11	12	26	43	40	44	36	38	29	18	21	13	333					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	14.3	67.4	70.6	139.8	249.2	246.3	317.4	320.4	402.3	328.9	233.5	365.6	254.0	138.9	10.8	94.7	127.1		
	P	0	0	0	0	0	7.3	34.5	36.4	70.9	125.9	123.3	157.7	159.0	204.2	171.8	129.1	227.3	175.6	70.5	5.5	48.3	65.9		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1					
	F	0	0	0	0	0	0	2	2	6	5	11	6	9	5	3	4	7	9	69					
	P	0	0	0	0	0	0	2	2	6	5	11	7	9	5	3	4	7	9	70					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	7.1	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4		
	F	0	0	0	0	0	0	12.3	11.8	32.3	29.0	67.7	43.3	80.1	52.9	34.0	51.9	121.9	175.8	28.8	1.8	17.7	25.3		
	P	0	0	0	0	0	0	6.3	6.1	16.4	14.6	33.9	25.1	39.8	26.9	17.8	28.7	75.8	121.6	14.8	0.9	9.3	13.7		
<b>C53. Cervix</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	1	3	2	3	3	0	2	2	2	3	0	2	2	25					
	P	0	0	0	0	0	1	3	2	3	3	0	2	2	2	3	0	2	2	25					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	F	0	0	0	0	0	7.2	18.4	11.8	16.1	17.4	0	14.4	17.8	21.2	34.0	0	34.8	39.1	10.4	0.8	7.4	9.8		
	P	0	0	0	0	0	3.6	9.4	6.1	8.2	8.8	0	7.2	8.8	10.7	17.8	0	21.7	27.0	5.3	0.4	3.8	5.1		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	2	1	1	4	1	2	0	0	0	1	3	15					
	P	0	0	0	0	0	0	0	2	1	1	4	1	2	0	0	0	1	3	15					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	F	0	0	0	0	0	0	11.8	5.4	5.8	24.6	7.2	17.8	0	0	0	17.4	58.6	6.3	0.4	4.0	5.6			
	P	0	0	0	0	0	0	6.1	2.7	2.9	12.3	3.6	8.8	0	0	0	10.8	40.5	3.2	0.2	2.1	3.0			

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C54, C55. Uterus</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	1	2	1	6	7	6	2	10	3	4	42					
	P	0	0	0	0	0	0	0	0	1	2	1	6	7	6	2	10	3	4	42					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	5.4	11.6	6.2	43.3	62.3	63.5	22.7	129.7	52.2	78.2	17.5	1.1	9.9	14.9		
	P	0	0	0	0	0	0	0	0	2.7	5.9	3.1	21.5	30.9	32.2	11.8	71.7	32.5	54.0	8.9	0.5	5.1	8.0		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	2	2	1	1	0	1	0	7						
	P	0	0	0	0	0	0	0	0	0	0	2	2	1	1	0	1	0	7						
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	14.4	17.8	10.6	11.3	0	17.4	0	2.9	0.3	1.9	2.5			
	P	0	0	0	0	0	0	0	0	0	0	7.2	8.8	5.4	5.9	0	10.8	0	1.5	0.1	1.0	1.3			
<b>C51, C52, C57. Vulva etc</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	1	3	2	2	2	1	0	1	12					
	P	0	0	0	0	0	0	0	0	0	0	1	3	2	2	2	1	0	1	12					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	6.2	21.6	17.8	21.2	22.7	13.0	0	19.5	5.0	0.4	3.2	4.4		
	P	0	0	0	0	0	0	0	0	0	0	3.1	10.8	8.8	10.7	11.8	7.2	0	13.5	2.5	0.2	1.6	2.3		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	2					
	P	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	2					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	0	0	0	0	0	19.5	0.8	0.0	0.4	0.6		
	P	0	0	0	0	0	0	0	0	0	0	0	3.6	0	0	0	0	0	13.5	0.4	0.0	0.2	0.4		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C56. Ovary</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	4	2	2	3	4	4	2	2	5	28				
	P	0	0	0	0	0	0	0	0	0	4	2	2	3	4	4	2	2	5	28				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	23.2	12.3	14.4	26.7	42.3	45.4	25.9	34.8	97.7	11.7	0.8	6.8	9.9	
	P	0	0	0	0	0	0	0	0	0	11.7	6.2	7.2	13.3	21.5	23.7	14.3	21.7	67.5	5.9	0.4	3.5	5.4	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	1	5	4	0	3	2	1	2	4	22				
	P	0	0	0	0	0	0	0	0	0	1	5	4	0	3	2	1	2	4	22				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	5.8	30.8	28.9	0	31.8	22.7	13.0	34.8	78.2	9.2	0.6	5.1	7.8	
	P	0	0	0	0	0	0	0	0	0	2.9	15.4	14.3	0	16.1	11.8	7.2	21.7	54.0	4.7	0.3	2.7	4.2	
<b>C58. Placenta</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C61. Prostate</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	2	7	27	41	57	65	44	40	29	312				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	2	7	27	41	57	65	44	40	29	312				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	11.8	43.2	192.3	359.6	622.3	806.2	705.7	1,145.1	1,268.6		133.9	10.2	78.9	133.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	5.9	21.6	96.8	181.1	306.4	385.1	315.5	433.0	391.7		66.0	5.0	36.7	59.1
Deaths																								
	M	0	0	0	0	0	0	0	0	0	2	2	3	7	10	12	14	20	25	95				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	2	2	3	7	10	12	14	20	25	95				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	11.8	12.3	21.4	61.4	109.2	148.8	224.5	572.6	1,093.6		40.8	1.8	21.5	44.8
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	5.9	6.2	10.8	30.9	53.7	71.1	100.4	216.5	337.7		20.1	0.9	9.1	17.9
<b>C62. Testis</b>																								
Cases																								
	M	0	0	0	0	1	1	1	5	7	1	1	1	0	0	1	0	0	0	19				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	1	1	1	5	7	1	1	1	0	0	1	0	0	0	19				
Incidence per 100,000																								
	M	0	0	0	0	6.9	7.4	6.4	31.2	38.7	5.9	6.2	7.1	0	0	12.4	0	0	0		8.2	0.6	6.9	8.4
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0
	P	0	0	0	0	3.5	3.6	3.1	15.1	19.1	2.9	3.1	3.6	0	0	5.9	0	0	0		4.0	0.3	3.4	4.1
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		0	0.0	0.0	0.0

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C60, C63. Penis etc</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	1	0	4				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	1	0	4				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	8.8	0	12.4	16.0	28.6	0	1.7	0.1	0.9	1.7	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	4.4	0	5.9	7.2	10.8	0	0.8	0.1	0.4	0.8	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	
<b>C64. Kidney</b>																								
Cases																								
	M	0	0	0	0	0	0	0	2	1	3	8	3	5	7	7	3	5	2	46				
	F	0	0	0	0	0	2	0	1	0	0	3	1	2	5	5	5	1	1	26				
	P	0	0	0	0	0	2	0	3	1	3	11	4	7	12	12	8	6	3	72				
Incidence per 100,000																								
	M	0	0	0	0	0	0	12.5	5.5	17.7	49.4	21.4	43.9	76.4	86.8	48.1	143.1	87.5	19.7	1.6	12.9	19.4		
	F	0	0	0	0	0	14.3	0	5.9	0	0	18.5	7.2	17.8	52.9	56.7	64.9	17.4	19.5	10.8	0.9	7.0	9.9	
	P	0	0	0	0	0	7.3	0	9.1	2.7	8.8	33.9	14.3	30.9	64.5	71.1	57.4	65.0	40.5	15.2	1.2	9.8	14.2	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	1	3	2	2	1	4	1	14					
	F	0	0	0	0	0	0	0	0	0	0	1	0	1	0	1	2	0	6					
	P	0	0	0	0	0	0	0	0	0	0	1	1	4	2	3	3	4	2	20				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	7.1	26.3	21.8	24.8	16.0	114.5	43.7	6.0	0.4	3.4	6.0		
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	8.9	0	11.3	25.9	0	2.5	0.1	1.2	2.1		
	P	0	0	0	0	0	0	0	0	0	0	3.1	3.6	17.7	10.7	17.8	21.5	43.3	4.2	0.3	2.2	3.8		



# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C67. Bladder</b>																									
Cases																									
	M	0	0	0	0	0	0	0	1	2	6	4	11	12	16	6	21	9	10	98					
	F	0	0	0	0	0	0	0	0	1	0	1	1	2	4	3	5	6	6	29					
	P	0	0	0	0	0	0	0	1	3	6	5	12	14	20	9	26	15	16	127					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	6.2	11.1	35.5	24.7	78.4	105.3	174.7	74.4	336.8	257.7	437.4	42.1	2.6	25.3	41.9		
	F	0	0	0	0	0	0	0	0	5.4	0	6.2	7.2	17.8	42.3	34.0	64.9	104.5	117.2	12.1	0.6	5.3	9.7		
	P	0	0	0	0	0	0	0	3.0	8.2	17.6	15.4	43.0	61.8	107.5	53.3	186.5	162.4	216.1	26.9	1.5	14.7	24.2		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	2	2	1	8	3	2	18					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	1	4					
	P	0	0	0	0	0	0	0	0	0	0	0	0	2	2	1	10	4	3	22					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	17.5	21.8	12.4	128.3	85.9	87.5	7.7	0.3	3.8	8.0		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	25.9	17.4	19.5	1.7	0.0	0.4	1.3		
	P	0	0	0	0	0	0	0	0	0	0	0	0	8.8	10.7	5.9	71.7	43.3	40.5	4.7	0.1	1.9	4.2		
<b>C65, C66, C68. Renal pelvis etc</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	2					
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1					
	P	0	0	0	0	0	0	0	0	0	0	0	0	2	1	0	0	0	0	3					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	8.8	10.9	0	0	0	0	0.9	0.1	0.7	0.8		
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	0	0	0	0	0	0.4	0.0	0.4	0.4		
	P	0	0	0	0	0	0	0	0	0	0	0	0	8.8	5.4	0	0	0	0	0.6	0.1	0.5	0.6		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2					
	F	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	1	4					
	P	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	2	1	6					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	10.9	0	0	28.6	0	0.9	0.1	0.5	0.9		
	F	0	0	0	0	0	0	0	0	0	0	0	0	8.9	0	0	13.0	17.4	19.5	1.7	0.0	0.7	1.3		
	P	0	0	0	0	0	0	0	0	0	0	0	0	4.4	5.4	0	7.2	21.7	13.5	1.3	0.0	0.6	1.1		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C69. Eye</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	2					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	28.6	0	0.4	0.0	0.1	0.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0.4	0.0	0.1	0.3		
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.2	10.8	0	0.4	0.0	0.1	0.4		
Deaths																									
	M	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1					
Mortality per 100,000																									
	M	0	0	0	0	0	0	6.4	0	0	0	0	0	0	0	0	0	0	0	0.4	0.0	0.4	0.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0		
	P	0	0	0	0	0	0	3.1	0	0	0	0	0	0	0	0	0	0	0	0.2	0.0	0.2	0.2		
<b>C71. Brain</b>																									
Cases																									
	M	0	2	0	0	0	0	3	0	2	1	4	3	1	2	3	1	1	0	23					
	F	0	0	0	0	0	1	1	0	0	1	4	0	2	2	2	0	1	0	14					
	P	0	2	0	0	0	1	4	0	2	2	8	3	3	4	5	1	2	0	37					
Incidence per 100,000																									
	M	0	11.8	0	0	0	0	19.3	0	11.1	5.9	24.7	21.4	8.8	21.8	37.2	16.0	28.6	0	9.9	0.8	7.5	9.6		
	F	0	0	0	0	0	7.2	6.1	0	0	5.8	24.6	0	17.8	21.2	22.7	0	17.4	0	5.8	0.5	4.4	5.6		
	P	0	6.1	0	0	0	3.6	12.6	0	5.5	5.9	24.7	10.8	13.3	21.5	29.6	7.2	21.7	0	7.8	0.7	5.9	7.5		
Deaths																									
	M	0	0	0	0	0	0	0	0	2	1	1	4	1	1	3	2	1	0	16					
	F	0	0	0	0	0	0	0	0	0	1	0	0	1	3	1	0	2	0	8					
	P	0	0	0	0	0	0	0	0	2	2	1	4	2	4	4	2	3	0	24					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	11.1	5.9	6.2	28.5	8.8	10.9	37.2	32.1	28.6	0	6.9	0.5	4.4	6.5		
	F	0	0	0	0	0	0	0	0	0	5.8	0	0	8.9	31.8	11.3	0	34.8	0	3.3	0.3	2.1	2.9		
	P	0	0	0	0	0	0	0	0	5.5	5.9	3.1	14.3	8.8	21.5	23.7	14.3	32.5	0	5.1	0.4	3.2	4.6		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C70, C72. Central nervous system</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0
<b>C73. Thyroid</b>																								
Cases																								
	M	0	1	0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	1	5				
	F	0	0	0	0	0	1	0	1	2	4	1	4	1	1	1	2	0	0	18				
	P	0	1	0	0	0	1	0	2	2	4	1	4	1	2	2	2	0	1	23				
Incidence per 100,000																								
	M	0	5.9	0	0	0	0	0	6.2	0	0	0	0	0	10.9	12.4	0	0	43.7	2.1	0.2	1.8	2.3	
	F	0	0	0	0	0	7.2	0	5.9	10.8	23.2	6.2	28.9	8.9	10.6	11.3	25.9	0	0	7.5	0.6	5.6	7.1	
	P	0	3.0	0	0	0	3.6	0	6.1	5.5	11.7	3.1	14.3	4.4	10.7	11.8	14.3	0	13.5	4.9	0.4	3.7	4.6	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	1	0	1	1	0	5				
	P	0	0	0	0	0	0	0	0	0	0	0	1	1	2	0	1	1	0	6				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	10.9	0	0	0	0	0.4	0.1	0.3	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	10.6	0	13.0	17.4	0	2.1	0.1	1.2	1.8	
	P	0	0	0	0	0	0	0	0	0	0	0	3.6	4.4	10.7	0	7.2	10.8	0	1.3	0.1	0.8	1.1	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C74, C75. Other endocrine</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	F	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1				
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	0	0	0	0	0	0	0	0.4	0.0	0.0	0.3	0.4
	P	0	0	0	0	0	0	0	0	0	0	3.1	0	0	0	0	0	0	0	0.2	0.0	0.0	0.2	0.2	
Deaths																									
	M	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1					
Mortality per 100,000																									
	M	0	0	0	0	0	0	6.4	0	0	0	0	0	0	0	0	0	0	0	0.4	0.0	0.0	0.4	0.5	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	3.1	0	0	0	0	0	0	0	0	0	0	0	0.2	0.0	0.0	0.2	0.2	
<b>C81. Hodgkins disease</b>																									
Cases																									
	M	0	0	1	1	1	0	1	0	1	0	0	1	0	0	0	0	0	0	6					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	2					
	P	0	0	1	1	1	0	1	0	1	0	0	1	0	1	0	1	0	0	8					
Incidence per 100,000																									
	M	0	0	5.7	5.8	6.9	0	6.4	0	5.5	0	0	7.1	0	0	0	0	0	0	2.6	0.2	2.6	2.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	0	13.0	0	0	0.8	0.1	0.4	0.7		
	P	0	0	2.9	2.9	3.5	0	3.1	0	2.7	0	0	3.6	0	5.4	0	7.2	0	0	1.7	0.1	1.5	1.7		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	0	0	0	0	0	0	0	0	0.4	0.0	0.4	0.4		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	2.9	0	0	0	0	0	0	0	0	0.2	0.0	0.2	0.2		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C82. Nodular NHL</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	2					
	F	0	0	0	0	0	0	1	0	1	1	1	3	1	0	3	0	0	1	12					
	P	0	0	0	0	0	0	1	0	1	1	2	3	1	0	4	0	0	1	14					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	6.2	0	0	0	12.4	0	0	0	0.9	0.1	0.6	0.8		
	F	0	0	0	0	0	0	6.1	0	5.4	5.8	6.2	21.6	8.9	0	34.0	0	0	19.5	5.0	0.4	3.3	4.6		
	P	0	0	0	0	0	0	3.1	0	2.7	2.9	6.2	10.8	4.4	0	23.7	0	0	13.5	3.0	0.3	2.0	2.8		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	2					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	3					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2	1	0	0	5					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	10.9	12.4	0	0	0	0.9	0.1	0.6	0.8		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	10.6	11.3	13.0	0	0	1.3	0.1	0.7	1.1		
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	10.7	11.8	7.2	0	0	1.1	0.1	0.6	1.0		
<b>C83. Diffuse NHL</b>																									
Cases																									
	M	0	1	0	0	2	0	0	1	2	2	1	4	5	1	8	0	3	2	32					
	F	0	0	0	0	0	0	0	0	1	1	0	2	2	4	2	3	1	2	18					
	P	0	1	0	0	2	0	0	1	3	3	1	6	7	5	10	3	4	4	50					
Incidence per 100,000																									
	M	0	5.9	0	0	13.7	0	0	6.2	11.1	11.8	6.2	28.5	43.9	10.9	99.2	0	85.9	87.5	13.7	1.2	9.8	13.5		
	F	0	0	0	0	0	0	0	0	5.4	5.8	0	14.4	17.8	42.3	22.7	38.9	17.4	39.1	7.5	0.5	4.4	6.4		
	P	0	3.0	0	0	7.0	0	0	3.0	8.2	8.8	3.1	21.5	30.9	26.9	59.2	21.5	43.3	54.0	10.6	0.9	7.0	9.7		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	1	1	2	2	2	3	2	2	0	15					
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	2	0	1	7					
	P	0	0	0	0	0	0	0	0	0	1	1	3	3	3	4	4	2	1	22					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	14.2	17.5	21.8	37.2	32.1	57.3	0	6.4	0.5	3.9	6.1		
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	10.6	11.3	25.9	0	19.5	2.9	0.2	1.5	2.5		
	P	0	0	0	0	0	0	0	0	0	2.9	3.1	10.8	13.3	16.1	23.7	28.7	21.7	13.5	4.7	0.3	2.7	4.2		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C84. T-cell lymphoma</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	1	0	1	0	0	1	1	0	0	0	4				
	F	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	2				
	P	0	0	0	0	0	0	0	0	1	0	1	1	0	1	1	1	0	0	6				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	5.5	0	6.2	0	0	10.9	12.4	0	0	0	1.7	0.2	1.2	1.6	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	0	0	0	13.0	0	0	0.8	0.0	0.4	0.7	
	P	0	0	0	0	0	0	0	0	2.7	0	3.1	3.6	0	5.4	5.9	7.2	0	0	1.3	0.1	0.8	1.2	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	3				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.4	0	0	0	0.4	0.1	0.2	0.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	19.5	0.8	0.0	0.2	0.6	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	7.2	0	13.5	0.6	0.0	0.3	0.6	
<b>C85. Other NHL</b>																								
Cases																								
	M	0	0	0	0	0	0	0	1	1	0	0	0	1	3	1	1	1	0	9				
	F	0	0	0	0	0	0	0	0	0	0	0	1	2	1	1	2	2	2	11				
	P	0	0	0	0	0	0	0	1	1	0	0	1	3	4	2	3	3	2	20				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	6.2	5.5	0	0	0	8.8	32.8	12.4	16.0	28.6	0	3.9	0.3	2.6	3.7	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	17.8	10.6	11.3	25.9	34.8	39.1	4.6	0.2	2.2	3.7	
	P	0	0	0	0	0	0	0	3.0	2.7	0	0	3.6	13.3	21.5	11.8	21.5	32.5	27.0	4.2	0.3	2.4	3.8	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	2	5				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2	1	3	7				
	P	0	0	0	0	0	0	0	0	0	0	0	0	1	0	2	3	1	5	12				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	8.8	0	12.4	16.0	0	87.5	2.1	0.1	1.2	2.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	11.3	25.9	17.4	58.6	2.9	0.1	0.9	2.2	
	P	0	0	0	0	0	0	0	0	0	0	0	0	4.4	0	11.8	21.5	10.8	67.5	2.5	0.1	1.0	2.3	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C82 - C85. Non-Hodgkin's lymphoma</b>																								
Cases																								
	M	0	1	0	0	2	0	0	2	4	2	3	4	6	5	11	1	4	2	47				
	F	0	0	0	0	0	0	1	0	2	2	1	7	5	5	6	6	3	5	43				
	P	0	1	0	0	2	0	1	2	6	4	4	11	11	10	17	7	7	7	90				
Incidence per 100,000																								
	M	0	5.9	0	0	13.7	0	0	12.5	22.1	11.8	18.5	28.5	52.6	54.6	136.4	16.0	114.5	87.5	20.2	1.8	14.2	19.7	
	F	0	0	0	0	0	0	6.1	0	10.8	11.6	6.2	50.5	44.5	52.9	68.1	77.8	52.2	97.7	17.9	1.3	10.3	15.4	
	P	0	3.0	0	0	7.0	0	3.1	6.1	16.4	11.7	12.3	39.4	48.6	53.7	100.7	50.2	75.8	94.5	19.0	1.5	12.2	17.5	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	1	1	2	3	3	6	3	2	2	23				
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	2	3	6	1	5	19				
	P	0	0	0	0	0	0	0	0	0	1	1	3	4	5	9	9	3	7	42				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	5.9	6.2	14.2	26.3	32.8	74.4	48.1	57.3	87.5	9.9	0.8	6.0	9.7	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	21.2	34.0	77.8	17.4	97.7	7.9	0.4	3.3	6.3	
	P	0	0	0	0	0	0	0	0	0	2.9	3.1	10.8	17.7	26.9	53.3	64.5	32.5	94.5	8.9	0.6	4.6	8.0	
<b>C81 - C85. All lymphomas</b>																								
Cases																								
	M	0	1	1	1	3	0	1	2	5	2	3	5	6	5	11	1	4	2	53				
	F	0	0	0	0	0	0	1	0	2	2	1	7	5	6	6	7	3	5	45				
	P	0	1	1	1	3	0	2	2	7	4	4	12	11	11	17	8	7	7	98				
Incidence per 100,000																								
	M	0	5.9	5.7	5.8	20.6	0	6.4	12.5	27.6	11.8	18.5	35.6	52.6	54.6	136.4	16.0	114.5	87.5	22.8	2.0	16.8	22.3	
	F	0	0	0	0	0	0	6.1	0	10.8	11.6	6.2	50.5	44.5	63.5	68.1	90.8	52.2	97.7	18.8	1.3	10.7	16.1	
	P	0	3.0	2.9	2.9	10.5	0	6.3	6.1	19.1	11.7	12.3	43.0	48.6	59.1	100.7	57.4	75.8	94.5	20.7	1.6	13.8	19.2	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	2	1	2	3	3	6	3	2	2	24				
	F	0	0	0	0	0	0	0	0	0	0	0	1	1	2	3	6	1	5	19				
	P	0	0	0	0	0	0	0	0	0	2	1	3	4	5	9	9	3	7	43				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	11.8	6.2	14.2	26.3	32.8	74.4	48.1	57.3	87.5	10.3	0.8	6.3	10.2	
	F	0	0	0	0	0	0	0	0	0	0	0	7.2	8.9	21.2	34.0	77.8	17.4	97.7	7.9	0.4	3.3	6.3	
	P	0	0	0	0	0	0	0	0	0	5.9	3.1	10.8	17.7	26.9	53.3	64.5	32.5	94.5	9.1	0.6	4.8	8.2	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C88. Immunoproliferative diseases</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	3	0	1	0	0	0	0	0	4				
	F	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	2				
	P	0	0	0	0	0	0	0	0	0	0	4	0	2	0	0	0	0	0	6				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	18.5	0	8.8	0	0	0	0	0	1.7	0.1	1.3	1.6	
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	8.9	0	0	0	0	0	0.8	0.1	0.7	0.8	
	P	0	0	0	0	0	0	0	0	0	0	12.3	0	8.8	0	0	0	0	0	1.3	0.1	1.0	1.2	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	43.7	0.4	0.0	0.2	0.6	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0.4	0.0	0.1	0.3	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.2	0	13.5	0.4	0.0	0.1	0.4	
<b>C90. Multiple myeloma</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	1	5	1	4	2	2	4	2	1	22				
	F	0	0	0	0	0	0	0	0	0	0	1	0	2	1	0	4	0	4	12				
	P	0	0	0	0	0	0	0	0	0	1	6	1	6	3	2	8	2	5	34				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	5.9	30.9	7.1	35.1	21.8	24.8	64.2	57.3	43.7	9.4	0.6	5.9	9.2	
	F	0	0	0	0	0	0	0	0	0	0	6.2	0	17.8	10.6	0	51.9	0	78.2	5.0	0.2	2.2	4.0	
	P	0	0	0	0	0	0	0	0	0	2.9	18.5	3.6	26.5	16.1	11.8	57.4	21.7	67.5	7.2	0.4	4.0	6.5	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	2	0	4				
	F	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	1	3	7				
	P	0	0	0	0	0	0	0	0	0	1	1	1	1	0	0	1	3	3	11				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	5.9	0	0	8.8	0	0	0	57.3	0	1.7	0.1	1.0	1.8	
	F	0	0	0	0	0	0	0	0	0	0	6.2	7.2	0	0	0	13.0	17.4	58.6	2.9	0.1	1.1	2.2	
	P	0	0	0	0	0	0	0	0	0	2.9	3.1	3.6	4.4	0	0	7.2	32.5	40.5	2.3	0.1	1.1	2.1	



# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C91. Lymphoid leukaemia</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	4	1	3	1	1	0	2	3	15				
	F	1	0	1	0	0	0	1	0	0	0	1	0	1	0	1	1	1	1	9				
	P	1	0	1	0	0	0	1	0	0	0	5	1	4	1	2	1	3	4	24				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	24.7	7.1	26.3	10.9	12.4	0	57.3	131.2	6.4	0.4	4.1	6.7	
	F	6.7	0	6.0	0	0	0	6.1	0	0	0	6.2	0	8.9	0	11.3	13.0	17.4	19.5	3.8	0.2	2.9	3.4	
	P	3.3	0	2.9	0	0	0	3.1	0	0	0	15.4	3.6	17.7	5.4	11.8	7.2	32.5	54.0	5.1	0.3	3.4	4.7	
Deaths																								
	M	0	1	1	0	0	0	1	0	0	0	0	0	1	0	0	1	1	1	7				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	3				
	P	0	1	1	0	0	0	1	0	0	0	0	0	1	0	0	1	3	2	10				
Mortality per 100,000																								
	M	0	5.9	5.7	0	0	0	6.4	0	0	0	0	0	8.8	0	0	16.0	28.6	43.7	3.0	0.1	2.4	3.2	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	34.8	19.5	1.3	0.0	0.3	0.9	
	P	0	3.0	2.9	0	0	0	3.1	0	0	0	0	0	4.4	0	0	7.2	32.5	27.0	2.1	0.1	1.3	2.0	
<b>C91.0. Acute lymphoid leukaemia</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1				
	F	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2				
	P	1	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	3				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	6.2	0	0	0	0	0	0	0	0.4	0.0	0.3	0.4	
	F	6.7	0	6.0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.8	0.1	1.3	0.9	
	P	3.3	0	2.9	0	0	0	0	0	0	0	3.1	0	0	0	0	0	0	0	0.6	0.0	0.8	0.6	
Deaths																								
	M	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	3				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
	P	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	3				
Mortality per 100,000																								
	M	0	5.9	5.7	0	0	0	6.4	0	0	0	0	0	0	0	0	0	0	0	1.3	0.1	1.5	1.3	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	3.0	2.9	0	0	0	3.1	0	0	0	0	0	0	0	0	0	0	0	0.6	0.0	0.8	0.7	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C91.1. Chronic lymphoid leukaemia</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	2	1	3	0	1	0	2	3	12					
	F	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	1	1	1	6					
	P	0	0	0	0	0	0	1	0	0	0	3	1	3	0	2	1	3	4	18					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	12.3	7.1	26.3	0	12.4	0	57.3	131.2	5.2	0.3	3.1	5.5		
	F	0	0	0	0	0	0	6.1	0	0	0	6.2	0	0	0	11.3	13.0	17.4	19.5	2.5	0.1	1.2	2.2		
	P	0	0	0	0	0	0	3.1	0	0	0	9.2	3.6	13.3	0	11.8	7.2	32.5	54.0	3.8	0.2	2.1	3.5		
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	3					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	2					
	P	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	2	2	5					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	8.8	0	0	0	28.6	43.7	1.3	0.0	0.7	1.5		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	17.4	19.5	0.8	0.0	0.2	0.6		
	P	0	0	0	0	0	0	0	0	0	0	0	0	4.4	0	0	0	21.7	27.0	1.1	0.0	0.4	0.9		
<b>C92. Myeloid leukaemia</b>																									
Cases																									
	M	1	1	0	0	0	0	1	0	0	0	1	0	2	1	4	2	1	2	16					
	F	0	1	0	0	0	0	0	0	1	0	0	0	0	0	3	2	3	4	14					
	P	1	2	0	0	0	0	1	0	1	0	1	0	2	1	7	4	4	6	30					
Incidence per 100,000																									
	M	6.4	5.9	0	0	0	0	6.4	0	0	0	6.2	0	17.5	10.9	49.6	32.1	28.6	87.5	6.9	0.5	5.0	7.0		
	F	0	6.2	0	0	0	0	0	0	5.4	0	0	0	0	0	34.0	25.9	52.2	78.2	5.8	0.2	2.5	4.6		
	P	3.3	6.1	0	0	0	0	3.1	0	2.7	0	3.1	0	8.8	5.4	41.5	28.7	43.3	81.0	6.3	0.4	3.8	5.8		
Deaths																									
	M	1	0	0	0	0	0	0	1	1	0	1	0	1	4	2	3	0	1	15					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	2	5	3	13					
	P	1	0	0	0	0	0	0	1	1	0	1	0	1	4	5	5	5	4	28					
Mortality per 100,000																									
	M	6.4	0	0	0	0	0	0	6.2	5.5	0	6.2	0	8.8	43.7	24.8	48.1	0	43.7	6.4	0.5	4.6	6.3		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	34.0	25.9	87.0	58.6	5.4	0.2	1.7	4.1		
	P	3.3	0	0	0	0	0	0	3.0	2.7	0	3.1	0	4.4	21.5	29.6	35.9	54.1	54.0	5.9	0.3	3.2	5.4		

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C92.0. Acute myeloid leukaemia</b>																								
Cases																								
	M	1	1	0	0	0	0	1	0	0	0	1	0	1	0	3	2	0	1	11				
	F	0	1	0	0	0	0	0	0	1	0	0	0	0	0	2	1	3	3	11				
	P	1	2	0	0	0	0	1	0	1	0	1	0	1	0	5	3	3	4	22				
Incidence per 100,000																								
	M	6.4	5.9	0	0	0	0	6.4	0	0	0	6.2	0	8.8	0	37.2	32.1	0	43.7	4.7	0.4	3.7	4.8	
	F	0	6.2	0	0	0	0	0	0	5.4	0	0	0	0	0	22.7	13.0	52.2	58.6	4.6	0.2	2.1	3.6	
	P	3.3	6.1	0	0	0	0	3.1	0	2.7	0	3.1	0	4.4	0	29.6	21.5	32.5	54.0	4.7	0.3	2.9	4.3	
Deaths																								
	M	1	0	0	0	0	0	1	0	0	1	0	1	3	2	2	2	0	1	12				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	1	4	3	11				
	P	1	0	0	0	0	0	0	1	0	0	1	0	1	3	5	3	4	4	23				
Mortality per 100,000																								
	M	6.4	0	0	0	0	0	6.2	0	0	6.2	0	8.8	32.8	24.8	32.1	0	43.7	5.2	0.4	3.8	5.1		
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	34.0	13.0	69.6	58.6	4.6	0.2	1.5	3.5		
	P	3.3	0	0	0	0	0	3.0	0	0	3.1	0	4.4	16.1	29.6	21.5	43.3	54.0	4.9	0.3	2.7	4.4		
<b>C92.1. Chronic myeloid leukaemia</b>																								
Cases																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	1	1	4				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2				
	P	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	1	2	6				
Incidence per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	8.8	10.9	0	0	28.6	43.7	1.7	0.1	1.0	1.8	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	19.5	0.8	0.0	0.2	0.6	
	P	0	0	0	0	0	0	0	0	0	0	0	0	4.4	5.4	0	7.2	10.8	27.0	1.3	0.0	0.6	1.1	
Deaths																								
	M	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	3				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1				
	P	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	2	0	0	4				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	5.5	0	0	0	0	10.9	0	16.0	0	0	1.3	0.1	0.8	1.2	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	13.0	0	0	0.4	0.0	0.1	0.3	
	P	0	0	0	0	0	0	0	0	2.7	0	0	0	0	5.4	0	14.3	0	0	0.8	0.0	0.5	0.8	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C93 - C95. Other &amp; unspecified leukaemia</b>																								
Cases																								
	M	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	4				
	F	0	0	0	0	0	1	0	0	0	0	0	0	0	0	2	0	0	0	3				
	P	0	1	1	0	0	1	0	0	0	0	0	0	0	0	2	2	0	0	7				
Incidence per 100,000																								
	M	0	5.9	5.7	0	0	0	0	0	0	0	0	0	0	0	0	32.1	0	0	1.7	0.1	1.4	1.7	
	F	0	0	0	0	0	7.2	0	0	0	0	0	0	0	0	22.7	0	0	0	1.3	0.1	1.0	1.3	
	P	0	3.0	2.9	0	0	3.6	0	0	0	0	0	0	0	0	11.8	14.3	0	0	1.5	0.1	1.2	1.5	
Deaths																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	2				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1				
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2	0	0	3				
Mortality per 100,000																								
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	32.1	0	0	0.9	0.0	0.3	0.9	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	11.3	0	0	0	0.4	0.1	0.2	0.4	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.9	14.3	0	0	0.6	0.0	0.3	0.6	
<b>C91 - C95. All leukaemia</b>																								
Cases																								
	M	1	1	1	0	0	0	1	0	0	0	5	1	5	2	5	4	3	5	34				
	F	1	1	1	0	0	1	1	0	1	0	1	0	1	0	6	3	4	5	26				
	P	2	2	2	0	0	1	2	0	1	0	6	1	6	2	11	7	7	10	60				
Incidence per 100,000																								
	M	6.4	5.9	5.7	0	0	0	6.4	0	0	0	30.9	7.1	43.9	21.8	62.0	64.2	85.9	218.7	14.6	1.0	9.9	15.0	
	F	6.7	6.2	6.0	0	0	7.2	6.1	0	5.4	0	6.2	0	8.9	0	68.1	38.9	69.6	97.7	10.8	0.6	6.5	9.3	
	P	6.5	6.1	5.9	0	0	3.6	6.3	0	2.7	0	18.5	3.6	26.5	10.7	65.2	50.2	75.8	135.1	12.7	0.8	8.1	11.8	
Deaths																								
	M	1	1	1	0	0	0	1	1	1	0	1	0	2	4	2	6	1	2	24				
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	2	7	4	17				
	P	1	1	1	0	0	0	1	1	1	0	1	0	2	4	6	8	8	6	41				
Mortality per 100,000																								
	M	6.4	5.9	5.7	0	0	0	6.4	6.2	5.5	0	6.2	0	17.5	43.7	24.8	96.2	28.6	87.5	10.3	0.6	7.3	10.4	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	45.4	25.9	121.9	78.2	7.1	0.2	2.2	5.3	
	P	3.3	3.0	2.9	0	0	0	3.1	3.0	2.7	0	3.1	0	8.8	21.5	35.5	57.4	86.6	81.0	8.7	0.4	4.8	7.9	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)		
<b>C80. Unspecified sites</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	3	1	2	5	2	5	8	14	4	5	49					
	F	0	0	0	0	0	1	0	0	0	1	1	0	1	1	0	8	4	13	30					
	P	0	0	0	0	0	1	0	0	3	2	3	5	3	6	8	22	8	18	79					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	16.6	5.9	12.3	35.6	17.5	54.6	99.2	224.5	114.5	218.7	21.0	1.2	11.6	21.2		
	F	0	0	0	0	0	7.2	0	0	0	5.8	6.2	0	8.9	10.6	0	103.8	69.6	254.0	12.5	0.2	4.6	9.5		
	P	0	0	0	0	0	3.6	0	0	8.2	5.9	9.2	17.9	13.3	32.2	47.4	157.8	86.6	243.1	16.7	0.7	8.0	15.1		
Deaths																									
	M	0	0	0	0	0	0	0	0	2	1	1	3	4	1	6	10	3	6	37					
	F	0	0	0	0	0	1	0	0	0	1	0	1	1	1	0	7	5	10	27					
	P	0	0	0	0	0	1	0	0	2	2	1	4	5	2	6	17	8	16	64					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	11.1	5.9	6.2	21.4	35.1	10.9	74.4	160.4	85.9	262.5	15.9	0.8	8.7	16.4		
	F	0	0	0	0	0	7.2	0	0	0	5.8	0	7.2	8.9	10.6	0	90.8	87.0	195.4	11.3	0.2	4.2	8.6		
	P	0	0	0	0	0	3.6	0	0	5.5	5.9	3.1	14.3	22.1	10.7	35.5	121.9	86.6	216.1	13.5	0.5	6.3	12.2		
<b>C96. Other haematopoietic</b>																									
Cases																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
Incidence per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
Deaths																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
Mortality per 100,000																									
	M	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	F	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	
	P	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	

# Cancer Incidence and Mortality in Tasmania 2002 by age, sex and site

		0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	Total	Crude Rate	Cumul Rate	AS Rates W AUS (1960) (2001)	
<b>C00 - C96. All cancers (excludes NMSCs)</b>																								
Cases																								
	<b>M</b>	2	6	2	2	6	7	12	18	38	40	92	127	164	194	256	206	132	94	1,398				
	<b>F</b>	1	1	2	1	2	17	21	28	50	92	85	99	109	137	139	138	112	102	1,136				
	<b>P</b>	3	7	4	3	8	24	33	46	88	132	177	226	273	331	395	344	244	196	2,534				
Incidence per 100,000																								
	<b>M</b>	12.7	35.3	11.5	11.5	41.2	51.8	77.2	112.4	210.0	236.6	567.7	904.8	1,438.6	2,117.9	3,175.0	3,303.9	3,779.0	4,112.0		600.1	45.0	374.4	592.1
	<b>F</b>	6.7	6.2	12.0	6.0	14.2	121.7	128.7	164.8	268.8	533.1	523.3	714.2	970.1	1,450.3	1,576.7	1,790.1	1,949.9	1,993.0		474.0	32.5	285.8	416.4
	<b>P</b>	9.8	21.2	11.7	8.8	27.9	87.4	103.6	139.3	239.8	386.4	545.5	810.1	1,206.0	1,779.0	2,340.2	2,467.0	2,641.6	2,647.2		536.2	38.6	325.7	491.9
Deaths																								
	<b>M</b>	1	1	2	1	1	1	3	3	10	15	25	45	62	78	114	123	77	77	639				
	<b>F</b>	0	0	0	1	0	1	4	7	12	13	32	41	47	42	54	75	75	85	489				
	<b>P</b>	1	1	2	2	1	2	7	10	22	28	57	86	109	120	168	198	152	162	1,128				
Mortality per 100,000																								
	<b>M</b>	6.4	5.9	11.5	5.8	6.9	7.4	19.3	18.7	55.3	88.7	154.3	320.6	543.9	851.5	1,413.9	1,972.7	2,204.4	3,368.3		274.3	17.5	158.7	279.1
	<b>F</b>	0	0	0	6.0	0	7.2	24.5	41.2	64.5	75.3	197.0	295.8	418.3	444.6	612.5	972.9	1,305.7	1,660.8		204.0	10.9	102.0	169.1
	<b>P</b>	3.3	3.0	5.9	5.9	3.5	7.3	22.0	30.3	60.0	82.0	175.7	308.3	481.5	645.0	995.3	1,420.0	1,645.6	2,188.0		238.7	14.1	127.3	215.7

# Coding differences between ICD-9 & ICD-10

*This table describes changes to the reporting of cancer sites and cancer groups from reporting in ICD-9 (in previous Cancer in Tasmania reports) to reporting in ICD-10 (in this report). Many of the changes have an effect on incidence and mortality rates.*

Reported in ICD-10 as:	Reported in earlier annual reports as:
Thymus, heart, mediastinum & pleura (ICD-10 C37, C38)	Previously reported as thymus, heart and mediastinum (ICD-9 164), but now includes non-mesothelioma pleura tumours (ICD-9 163).
Skin (C44)	Now includes Merkel cell carcinoma (ICD-9 8247/3).
Mesothelioma (ICD-10 C45)	Previously included in pleura (ICD-9 163) and peritoneum (ICD-9 158.8).
Kaposi sarcoma (ICD-10 C46)	Previously reported as soft tissue (ICD-9 171) and specified internal organs.
Connective tissue (ICD-10 C47, C49)	Previously reported as soft tissue (ICD-9 171) but no longer includes merkel cell carcinoma (ICD-9 8247/3) and Kaposi sarcoma (ICD-9 9140/3).
Vulva etc (ICD-10 C51, C52, C57)	Previously reported as vagina and vulva (ICD-9 184) and now includes other uterine adnexae (ICD-9 183.2 – 183.9).
Uterus (ICD-10 C54-C55)	Previously reported as corpus uteri (ICD-9 182).
Ovary (ICD-10 C56)	Previously reported as ovary (ICD-9 183) and included other uterine adnexae (ICD-9 183.2 – 183.9).
Kidney (ICD-10 C64)	Previously included in kidney (ICD-9 189) and included renal pelvis, ureter, urethra and other/unspecified urinary organs.
Renal pelvis etc (ICD-10 C65, C66, C68)	Previously included in kidney (ICD-9 189).
Central nervous system (ICD-10 C70, C72)	Previously reported as other central nervous system (ICD-9 192).
T-cell lymphoma (ICD-10 C84)	Previously included in all lymphomas and other lymphomas, and now only includes ICD-9 codes 202.1 and 202.2.
Other non-Hodgkin's lymphomas (ICD-10 C85).	Previously included in other lymphomas (ICD-9 202.1, 202.2, 202.8, 202.9) but now only includes unspecified non-Hodgkin's lymphoma (ICD-9 202.8).
Immunoproliferative diseases (ICD-10 C88)	Previously included in multiple myeloma (ICD-9 203).
Multiple myeloma (ICD-10 C90)	Previously reported as multiple myeloma (ICD-9 203), but no longer includes immunoproliferative neoplasms.
Other & unspecified haematopoietic neoplasms (ICD-10 C96)	Previously included in tumours of histiocytic tissue (ICD-9 202.3, 202.5 & 202.6) and in other lymphomas (under ICD-9 202.9).

---

# Appendices

---

Appendix A: Cancer site codes and combinations

Appendix B: Statistical methods

Appendix C: Population data

Appendix D: Indices of data quality

Appendix E: Use of Tasmanian Cancer Registry data

Appendix F: Incidence and mortality summary tables 2002



## Appendix A: Cancer site codes and combinations

*In past annual Cancer in Tasmania reports, cancers were reported in ICD-9 codes, however, in this report cancers have been reported according to ICD-10 codes. ICD-9 and ICD-10 codes are comparable for most cancers. Where these codes are not comparable an explanation of the differences between them is provided on page 54 Coding differences between ICD-9 & ICD-10.*

ICD-10 description	ICD-10
<b>Lip, oral cavity and pharynx (C00 - C14)</b>	
Lip	C00
Tongue	C01-C02
Gum	C03
Floor of mouth	C04
Other mouth	C05-C06
Oral cavity	C01-C06
Salivary glands	C07-C08
Oropharynx	C09-C10
Nasopharynx	C11
Hypopharynx (including pyriform sinus)	C12-C13
Pharynx	C09-C13
Other oral (includes other & unspecified sites of lip, oral cavity & pharynx)	C14
Head and neck	C01-C14, C30-C32
<b>Digestive organs (C15 - C26)</b>	
Oesophagus	C15
Stomach	C16
Small intestine (including duodenum)	C17
Colon	C18
Rectum (including rectosigmoid, anal canal & anus)	C19-C21
Colorectal	C18-C21
Liver (and intrahepatic bile ducts)	C22
Gallbladder (and other biliary tract)	C23-C24
Pancreas	C25
<b>Respiratory system and intrathoracic organs (C30 - C39)</b>	
Nasal cavities	C30-C31
Larynx	C32
Lung (includes trachea, bronchus & lung)	C33-C34
Thymus (includes heart, mediastinum & pleura)	C37-C38
<b>Bones, joints and articular cartilage (C40 - C41)</b>	
Bone (includes articular cartilage)	C40-C41
<b>Skin (C43, C44)</b>	
Melanoma of skin	C43
Skin (excludes melanoma)	C44
<b>Mesothelioma and connective tissue (C45 - C49)</b>	
Mesothelioma	C45
Kaposi sarcoma	C46
Retroperitoneum & peritoneum	C48
Connective tissue (includes peripheral nerves etc.)	C47, C49
<b>Breast (C50) and female genital organs (C51 - C58)</b>	
Breast	C50
Cervix	C53
Uterus	C54, C55
Ovary	C56
Placenta	C58
Vulva and other/unspecified female genital organs	C51, C52, C57

ICD-10 description	ICD-10
<b>Male genital organs (C60 - C63)</b>	
Prostate	C61
Testis	C62
Penis (and other male genital organs)	C60, C63
<b>Urinary tract (C64 - C68)</b>	
Kidney (except renal pelvis)	C64
Bladder	C67
Renal pelvis (including other/unspecified urinary organs)	C65, C66, C68
<b>Eye, brain and other parts of the central nervous system (C69 - C71)</b>	
Eye	C69
Brain	C71
Central nervous system (includes meninges)	C70, C72
<b>Thyroid and other endocrine glands (C73 - C75)</b>	
Thyroid	C73
Other endocrine (glands and related structures)	C74, C75
<b>Malignant neoplasms of lymphoid, haematopoietic and related tissue (C81 - C96)</b>	
Hodgkin's disease	C81
Nodular non-Hodgkin's lymphoma	C82
Diffuse non-Hodgkin's lymphoma	C83
T-cell lymphoma	C84
Other non-Hodgkin's lymphoma	C85
Non-Hodgkin's lymphoma	C82-C85
All lymphomas	C81-C85
Immunoproliferative disease	C88
Multiple myeloma (and malignant plasma cell neoplasms)	C90
Lymphoid leukaemia	C91
Acute lymphoid leukaemia	C91.0
Chronic lymphoid leukaemia	C91.1
Myeloid leukaemia	C92
Acute myeloid leukaemia	C92.0
Chronic myeloid leukaemia	C92.1
Other and unspecified leukaemia	C93 - C95
All leukaemia	C91-C95
Other haematopoietic neoplasms	C96
<b>Unknown primary site (C80)</b>	
Other and unspecified sites	C80
<b>All cancers (excluding non-melanoma skin cancers)</b>	<b>C00-C96</b>

---

## Appendix B: Statistical methods

---

The 2002 report contains numbers of new cases and deaths, and age specific, crude, cumulative, and age standardised incidence and mortality rates of Tasmanian residents diagnosed with cancer. They are based on registrations completed by 30 September 2004.

### Incidence

Cancer incidence is defined as the number of new cases of cancer in a population during a specific period. The incidence data in this report refer to the number of primary cancers first diagnosed between 1 January 2002 and 31 December 2002 in persons who were residents of Tasmania at the time of diagnosis.

### Mortality

The mortality data in this report refer to deaths from cancer of people who were first diagnosed as having cancer while they were residents of Tasmania. In many instances, other State or Territory cancer registries notify the Tasmanian Cancer Registry of cancer patients who die outside Tasmania. Details of patients diagnosed interstate who die in Tasmania are forwarded to the relevant cancer registry. Deaths from other causes are also recorded so that survival proportions can be calculated.

### Crude rates (CR)

The crude incidence (rate) is calculated as the number of new cases of cancer divided by the population at risk in a specified time period. The crude mortality rate substitutes deaths for new cases in this calculation. Both are conventionally expressed as annual rates per 100,000 population. The Australian Bureau of Statistics (ABS) supplied the estimated Tasmanian population by age and sex for 2002 in June 2003.

### Age specific rates

Age specific rates are calculated by dividing the number of cases occurring in each specified age group by the corresponding population in the same age group and are expressed as an annual rate per 100,000 population.

### Age standardised rates (ASR)

Rates are adjusted for age to facilitate comparisons between populations that have different age structures, e.g. between youthful and aging communities. In this publication we use direct standardisation in which age specific rates are used to calculate the number of cases that would have occurred if the population had the same age distribution as the World Standard Population 1960, World Standard Population 2000 and the Australian Standard Population 2001. This effectively removes the influence of age structure on the summary rate, which is described as the age standardised rate. The method may be used for both incidence and mortality calculations.

### Cumulative rates

The cumulative rate is a directly standardised rate calculated by summing the age specific rate for each year of life prior to age 75. Cumulative risk to age 75 can be calculated from the cumulative rate.

## Appendix C: Population data

### Estimated resident population of Tasmania by age group and sex for 30 June 2002<sup>1</sup>

Age	Males	Females	Persons
0-4	15,699	14,878	30,577
5-9	16,988	16,064	33,052
10-14	17,415	16,673	34,088
15-19	17,346	16,639	33,985
20-24	14,549	14,105	28,654
25-29	13,503	13,970	27,473
30-34	15,547	16,314	31,861
35-39	16,019	16,992	33,011
40-44	18,097	18,600	36,697
45-49	16,905	17,258	34,163
50-54	16,205	16,242	32,447
55-59	14,037	13,861	27,898
60-64	11,400	11,236	22,636
65-69	9,160	9,446	18,606
70-74	8,063	8,816	16,879
75-79	6,235	7,709	13,944
80-84	3,493	5,744	9,237
85+	2,286	5,118	7,404
<b>Total</b>	<b>232,947</b>	<b>239,665</b>	<b>472,612</b>

### Australian Standard Population and World Standard Populations

Age	Australian Standard Population (2001) <sup>2</sup>	World Standard Population (1960) <sup>3</sup>	World Standard Population (2000) <sup>4</sup>
0-4	1,282,357	12,000	8.86
5-9	1,351,664	10,000	8.69
10-14	1,353,177	9,000	8.60
15-19	1,352,745	9,000	8.47
20-24	1,302,412	8,000	8.22
25-29	1,407,081	8,000	7.93
30-34	1,466,615	6,000	7.61
35-39	1,492,204	6,000	7.15
40-44	1,479,257	6,000	6.59
45-49	1,358,594	6,000	6.04
50-54	1,300,777	5,000	5.37
55-59	1,008,799	4,000	4.55
60-64	822,024	4,000	3.72
65-69	682,513	3,000	2.96
70-74	638,380	2,000	2.21
75-79	519,356	1,000	1.52
80-84	330,050	500	0.91
85+	265,235	500	0.63
<b>Total</b>	<b>19,413,240</b>	<b>100,000</b>	<b>100.03</b>

The Australian Standard Population 2001<sup>2</sup> approximates the proportional age distribution of the whole of Australia in 2001. Similarly the World Standard Population<sup>3</sup> approximates the proportional age distribution of the whole world. Standard populations are used widely for direct standardisation to enable comparisons between populations of differing age structure.

<sup>1</sup> Australian Bureau of Statistics (ABS), Population by Age and Sex, Australian States and Territories, June 2003 (Cat. No. 3201.0).

<sup>2</sup> Australian Bureau of Statistics (ABS) 2004, Australian demographic statistics, March quarter 2004 (Cat. No. 3101.0).

<sup>3</sup> Parkin D M, Whelan S, Ferlay J, Raymond L and Young J. Cancer Incidence in Five Continents Vol VII, IARC Scientific Publications No. 143. Lyon: International Agency for Research on Cancer, 1997 (p 67).

<sup>4</sup> Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJL, Lozano R & Inoue M, 2000. Age standardization of rates: a new WHO standard. GPE Discussion Paper Series No. 31. Geneva: World Health Organisation (*available at* [http://www.who.int/health-systems-performance/docs/levelofhealth\\_docs.htm](http://www.who.int/health-systems-performance/docs/levelofhealth_docs.htm)).

## Appendix D: Indices of data quality

Three indices of data quality are commonly used by Australian Cancer Registries: the mortality to incidence ratio (M/I%), the proportion of cancers with histological verification (HV%) and the proportion of cancers registered on the basis of death certificate only (DCO%). The Tasmanian Cancer Registry has calculated these three indices and also determined the proportion of cancers with morphological verification (MV%) and the proportion of cancers of unknown primary site (PSU) for 2002 data.

### Death Certificate Only (DCO)

In the past, the Tasmanian Cancer Registry did not register cases on the basis of DCO, unlike other State and Territory cancer registries that registered these cases and included them in their reports. Each death certificate notification is actively followed up until the time and place of diagnosis are ascertained and the diagnosis verified. If the diagnostic details cannot be confirmed morphologically, the case is registered on the basis of a clinical diagnosis<sup>8</sup>. In 2002, 28 DCO cases were registered and these cases have been included in the 2002 incidence data. This will increase the number of new cancers by 1.1% (1.1% for both males and females). For DCO cases, the date of diagnosis is taken as the date of death. Where there is a low DCO%, as is the case for this Registry, the potential error in registration is reduced.

### Mortality to Incidence Ratio (M/I%)

One way of assessing the completeness of cancer ascertainment is the M/I%. This measure is calculated by dividing the number of deaths attributed to a specific cancer in a defined population by the number of new cases of the same cancer registered during the same period in the same population. For cancers with a poor prognosis, the ratio will be close to 100%. If it exceeds 100% this may indicate that the cancer is being under-registered, but a more likely explanation for this result with uncommon cancers is that it is a result of random fluctuations in the annual number of new cases and deaths.

---

<sup>8</sup> Note: Prior to 2000, DCO cases were not registered by the Tasmanian Cancer Registry.

### Histological Verification (HV%)

HV% is the percentage of cases with verification by histological investigation. Histological verification of diagnosis shows that it has been possible to investigate a patient with such thoroughness that a portion of the suspected neoplasm has been removed for microscopic examination<sup>9</sup>. For 2002, 81% of all male and 85% of all female registered cases had a diagnosis on the basis of tissue examination. HV% includes only the cancers that were diagnosed following tissue or needle biopsy and did not include diagnoses made on the basis of cytology examination of smears or aspirates (including haematological examinations).

### Morphological Verification (MV%)

If we add the diagnoses based on exfoliative cytology, haematological examinations (for leukaemia) as well as diagnoses based on histological examination of a tissue specimen, then the percentage of all cancers with morphological verification (MV%) in 2002 was 91% for males and 92% for females<sup>10</sup>.

### Unknown Primary Site (PSU)

The Registry calculated the percentage of all cancers that were classified as PSU (reported as ICD-10 C80 Unspecified site), because it is one of the quality indicators used for international comparisons<sup>11</sup>. In 2002, the percentage of all cases classified as PSU was 3.1% (3.5% for males and 2.6% for females).

---

<sup>9</sup> Parkin D.M, Chen V.W, Ferlay J, Galceran J, Storm H.H and Whelan S. *Comparability and Quality Control in Cancer Registration*. IARC Technical Report No 19. Lyon: International Agency for Research on Cancer, 1994 (p 43).

<sup>10</sup> Parkin D.M, Whelan S, Ferlay J, Raymond L and Young J. *Cancer Incidence in Five Continents Vol VII*. IARC Scientific Publication No 143. Lyon: International Agency for Research on Cancer, 1997 (p 50).

<sup>11</sup> Parkin D.M, Chen V.W, Ferlay J, Galceran J, Storm H.H and Whelan S. *Comparability and Quality Control in Cancer Registration*. IARC Technical Report No 19. Lyon: International Agency for Research on Cancer, 1994 (p 51).

**Table 2: Indices of Data Quality**

ICD-10	Site	Incidence	Mortality	M/I%	HV%	MV%	DCO%
<b>Males</b>							
C00	Lip	26	0	0	100	100	0
C01-C14,C30-C32	Head and neck	46	16	35	96	100	0
C15	Oesophagus	29	31	107	97	97	0
C16	Stomach	32	22	69	97	97	0
C18	Colon	118	46	39	94	94	0
C19 - C21	Rectum	76	31	41	99	99	0
C22	Liver	9	8	89	33	67	11
C23, C24	Gallbladder	12	10	83	75	75	8
C25	Pancreas	27	33	122	41	63	4
C33, C34	Lung	199	152	76	51	83	1
C43	Melanoma of skin	117	24	21	100	100	0
C50	Breast	0	1	-	-	-	-
C61	Prostate	312	95	30	82	93	2
C62	Testis	19	0	0	100	100	0
C64	Kidney	46	14	30	85	87	0
C67	Bladder	98	18	18	97	100	0
C71	Brain	23	16	70	74	74	0
C73	Thyroid	5	1	20	60	80	0
C81 - C85	All lymphomas	53	24	45	96	100	0
C90	Multiple myeloma	22	4	18	82	86	5
C91 - C95	All leukaemia	34	24	71	41	76	3
C80	Unspecified sites	49	37	76	45	57	6
<b>C00 - C96</b>	<b>All cancers (excludes NMSCs)</b>	<b>1398</b>	<b>639</b>	<b>46</b>	<b>81</b>	<b>91</b>	<b>1</b>
<b>Females</b>							
C00	Lip	12	0	0	100	100	0
C01-C14,C30-C32	Head and neck	8	5	63	100	100	0
C15	Oesophagus	12	7	58	92	92	0
C16	Stomach	21	17	81	95	95	0
C18	Colon	109	49	45	93	93	1
C19 - C21	Rectum	40	37	93	100	100	0
C22	Liver	7	4	57	14	14	14
C23, C24	Gallbladder	10	7	70	60	60	0
C25	Pancreas	37	25	68	49	54	5
C33, C34	Lung	105	112	107	49	78	1
C43	Melanoma of skin	108	5	5	100	100	0
C50	Breast	333	69	21	97	99	1
C53	Cervix	25	15	60	96	96	4
C54, C55	Uterus	42	7	17	98	98	0
C51, C52, C57	Vulva etc	12	2	17	100	100	0
C56	Ovary	28	22	79	79	96	0
C64	Kidney	26	6	23	85	88	0
C67	Bladder	29	4	14	90	90	0
C71	Brain	14	8	57	100	100	0
C73	Thyroid	18	5	28	94	100	0
C81 - C85	All lymphomas	45	19	42	84	98	0
C90	Multiple myeloma	12	7	58	67	83	17
C91 - C95	All leukaemia	26	17	65	42	92	0
C80	Unspecified sites	30	27	90	50	67	3
<b>C00 - C96</b>	<b>All cancers (excludes NMSCs)</b>	<b>1136</b>	<b>489</b>	<b>43</b>	<b>85</b>	<b>92</b>	<b>1</b>

# Appendix E: Use of Tasmanian Cancer Registry Data

## Confidentiality of information

Confidentiality of data is a requirement of the *Public Health Act 1997*. The Registry cannot release data identifying an individual unless authorised by the Director of Public Health.

The relevant sections of this Act are reproduced below.

### Disclosure of information relating to a notifiable disease

Section 61: A person, unless authorised to do so under section 147, must not disclose any information in relation to:

- (a) Any notification relating to a notifiable disease; or
- (b) Any investigation or inquiry into a notifiable disease; or
- (c) The identity of any person to whom any notification, investigation or inquiry relates.

### Disclosure of information

Section 147: A person must not disclose any information obtained for the purpose of this Act relating to a person except in accordance with any relevant guidelines and:

- (a) With the written consent of the person or parent or guardian of a child or person to whom the information relates; or
- (b) To a registered medical practitioner who is directly involved in the treatment of that person; or
- (c) To a person apparently in charge of any institution or facility which is involved in the diagnosis or treatment of that person; or
- (d) To a person authorised by the Director; or
- (e) For the purpose of notifying a notifiable disease; or
- (f) For the purpose of an epidemiological study or research authorised by the Director; or
- (g) For the purpose of legal proceedings arising out of this Act; or
- (h) For a purpose authorised or required by this Act or another Act; or
- (i) For the purposes of study or research approved by the Director.

## Requests for non-identifying data

Non-identifying cancer data are available upon request. Data are usually released as incidence or mortality rates, or number of cases or deaths, for specific cancers, cancer morphologies, time periods or age groups. Data that can be compiled using existing reporting systems have a turnaround of approximately 48 hours. Data needing input from the biostatistician require more notice. Such data are provided to epidemiological and clinical researchers, the Department of Health and Human Services, students and the public. It is generally not feasible to release data for small geographical areas, because this could lead to identification of the persons diagnosed and the Registry may not have population data for the area with which to assess the rate of occurrence. In addition to data requests, the Registry receives personal enquiries regarding cancer. When appropriate, these enquiries are referred to other agencies or health professionals.

## Requests for identifying data

The release of named data is strictly controlled. Named data may be released only after approval of a formal application submitted to the Director of the Tasmanian Cancer Registry, and with subsequent approval by the Director of Public Health. Applications for research purposes need the approval of the researchers' institutional ethics committee and the Southern Tasmania Health and Medical Human Research Ethics Committee. Prior to contacting persons named, researchers must notify the patient's medical practitioner of their intent to contact and provide a form to be returned if he/she does not want the patient to be approached.

Applicants are required to send a covering letter (detailing the purposes of the study and who is funding the study), copies of the study protocol (including the approach letter to participants) and two copies of the Ethics Committees approvals to the Director of the Tasmanian Cancer Registry.

## **Published data**

Annual reports from the Registry provide data on cancer numbers and incidence and mortality. Additional information is provided on selected cancer sites. It should be recognised that active follow-up is necessary to complete registrations for up to 30% of cases each year, which results in a two-year interval from year of diagnosis to date of publication of incidence data. Considerable time is spent on matching, classifying and validating cancer cases notified to the Registry. In addition the Tasmanian Cancer Registry supplies data to the National Cancer Statistics Clearing House (NCSCH) and to the International Association of Cancer Registries (IACR).



# Appendix F: Incidence and mortality summary tables 2002

Table 3: Incidence summary table 2002

ICD-10	Site	Males					Females					Total N
		*N	CR	ASRW 1960	ASRW 2000	ASRA 2001	N	CR	ASRW 1960	ASRW 2000	ASRA 2001	
C00	Lip	26	11.2	7.1	8.3	11.5	12	5.0	2.4	2.8	4.1	38
C01, C02	Tongue	12	5.2	3.3	3.7	4.8	1	0.4	0.3	0.4	0.4	13
C03	Gum	0	0.0	0.0	0.0	0.0	1	0.4	0.3	0.3	0.4	1
C04	Floor of mouth	3	1.3	0.8	0.9	1.2	0	0.0	0.0	0.0	0.0	3
C05, C06	Other mouth	3	1.3	0.9	0.9	1.2	2	0.8	0.5	0.6	0.8	5
C07, C08	Salivary glands	1	0.4	0.2	0.3	0.4	1	0.4	0.3	0.3	0.4	2
C09, C10	Oropharynx	4	1.7	1.2	1.3	1.6	0	0.0	0.0	0.0	0.0	4
C11	Nasopharynx	0	0	0	0	0	0	0	0	0	0	0
C12, C13	Hypopharynx	2	0.9	0.4	0.5	0.9	0	0.0	0.0	0.0	0.0	2
C14	Other oral	1	0.4	0.2	0.2	0.4	0	0.0	0.0	0.0	0.0	1
C15	Oesophagus	29	12.4	7.3	8.4	11.8	12	5.0	1.7	2.3	3.9	41
C16	Stomach	32	13.7	7.6	9.0	13.6	21	8.8	4.5	5.4	7.5	53
C17	Small intestine	8	3.4	2.4	2.6	3.7	5	2.1	1.3	1.6	1.9	13
C18	Colon	118	50.7	29.3	34.0	50.4	109	45.5	23.4	26.5	38.1	227
C19 - C21	Rectum	76	32.6	21.0	23.1	31.4	40	16.7	8.1	9.6	13.9	116
C18 - C21	Colorectal	194	83.3	50.3	57.1	81.8	149	62.2	31.5	36.1	52.0	343
C22	Liver	9	3.9	2.9	3.0	3.7	7	2.9	1.8	1.9	2.5	16
C23, C24	Gallbladder	12	5.2	3.1	3.5	5.3	10	4.2	1.9	2.1	3.2	22
C25	Pancreas	27	11.6	6.7	7.8	11.4	37	15.4	7.8	8.9	12.8	64
C30, C31	Nasal cavities	8	3.4	2.2	2.4	3.4	0	0.0	0.0	0.0	0.0	8
C32	Larynx	12	5.2	3.5	3.7	4.7	3	1.3	0.7	0.8	1.0	15
C33, C34	Lung	199	85.4	50.1	57.8	83.9	105	43.8	24.6	27.5	37.7	304
C37, C38	Thymus etc	1	0.4	0.3	0.3	0.4	1	0.4	0.1	0.2	0.3	2
C40, C41	Bone	3	1.3	1.3	1.2	1.2	1	0.4	0.4	0.3	0.4	4
C43	Melanoma of skin	117	50.2	36.1	39.7	49.1	108	45.1	32.2	34.9	42.2	225
C44	Skin	Incidence of non-melanoma skin cancer is not routinely registered.										
C45	Mesothelioma	12	5.2	3.1	3.6	4.8	1	0.4	0.3	0.3	0.4	13
C47, C49	Connective tissue	11	4.7	3.2	3.4	4.5	14	5.8	3.6	4.0	5.3	25
C50	Breast	0	0.0	0.0	0.0	0.0	333	138.9	94.7	102.8	127.1	333
C53	Cervix	-	-	-	-	-	25	10.4	7.4	8.2	9.8	25
C54, C55	Uterus	-	-	-	-	-	42	17.5	9.9	11.0	14.9	42
C51, C52, C57	Vulva etc	-	-	-	-	-	12	5.0	3.2	3.4	4.4	12
C56	Ovary	-	-	-	-	-	28	11.7	6.8	7.3	9.9	28
C58	Placenta	-	-	-	-	-	0	0	0	0	0	0
C61	Prostate	312	133.9	78.9	90.5	133.0	-	-	-	-	-	312
C62	Testis	19	8.2	6.9	7.7	8.4	-	-	-	-	-	19
C60, C63	Penis etc	4	1.7	0.9	1.1	1.7	-	-	-	-	-	4
C64	Kidney	46	19.7	12.9	14.3	19.4	26	10.8	7.0	7.6	9.9	72
C67	Bladder	98	42.1	25.3	29.2	41.9	29	12.1	5.3	6.4	9.7	127
C65, C66, C68	Renal pelvis etc	2	0.9	0.7	0.6	0.8	1	0.4	0.4	0.3	0.4	3
C69	Eye	1	0.4	0.1	0.3	0.5	1	0.4	0.1	0.2	0.3	2
C71	Brain	23	9.9	7.5	8.2	9.6	14	5.8	4.4	4.7	5.6	37
C70, C72	Central nervous system	0	0	0	0	0	0	0	0	0	0	0
C73	Thyroid	5	2.1	1.8	1.8	2.3	18	7.5	5.6	6.0	7.1	23
C74, C75	Other endocrine	0	0.0	0.0	0.0	0.0	1	0.4	0.3	0.3	0.4	1
C81 - C85	All lymphomas	53	22.8	16.8	18.1	22.3	45	18.8	10.7	12.0	16.1	98
C90	Multiple myeloma	22	9.4	5.9	6.6	9.2	12	5.0	2.2	2.6	4.0	34
C91 - C95	All leukaemia	34	14.6	9.9	10.8	15.0	26	10.8	6.5	7.0	9.3	60
C80	Unspecified sites	49	21.0	11.6	14.0	21.2	30	12.5	4.6	5.7	9.5	79
<b>C00 - C96</b>	<b>All cancers (excl. NMSC)</b>	<b>1398</b>	<b>600.1</b>	<b>374.4</b>	<b>424.2</b>	<b>592.1</b>	<b>1136</b>	<b>474.0</b>	<b>285.8</b>	<b>317.0</b>	<b>416.4</b>	<b>2534</b>

**Table 4: Mortality summary table 2002**

ICD-10	Site	Males					N	CR	Females			Total N
		*N	CR	ASW 1960	ASRW 2000	ASRA 2001			ASRW 1960	ASRW 2000	ASRA 2001	
C00	Lip	0	0	0	0	0	0	0	0	0	0	0
C01, C02	Tongue	6	2.6	1.7	1.9	2.6	0	0.0	0.0	0.0	0.0	6
C03	Gum	0	0	0	0	0	0	0	0	0	0	0
C04	Floor of mouth	1	0.4	0.2	0.3	0.4	0	0.0	0.0	0.0	0.0	1
C05, C06	Other mouth	2	0.9	0.5	0.6	0.8	1	0.4	0.3	0.3	0.4	3
C07, C08	Salivary glands	1	0.4	0.2	0.3	0.6	0	0.0	0.0	0.0	0.0	1
C09, C10	Oropharynx	0	0.0	0.0	0.0	0.0	2	0.8	0.5	0.6	0.8	2
C11	Nasopharynx	0	0.0	0.0	0.0	0.0	1	0.4	0.1	0.1	0.3	1
C12,C13	Hypopharynx	1	0.4	0.2	0.3	0.4	1	0.4	0.1	0.2	0.3	2
C14	Other oral	0	0	0	0	0	0	0	0	0	0	0
C15	Oesophagus	31	13.3	8.0	9.1	12.8	7	2.9	1.5	1.6	2.3	38
C16	Stomach	22	9.4	5.2	6.2	10.0	17	7.1	4.0	4.6	6.2	39
C17	Small intestine	4	1.7	0.8	1.1	2.1	1	0.4	0.1	0.2	0.3	5
C18	Colon	46	19.7	11.3	13.1	20.3	49	20.4	8.9	10.8	16.3	95
C19 - C21	Rectum	31	13.3	7.5	8.9	14.1	37	15.4	6.7	8.1	12.4	68
C18 - C21	Colorectal	77	33.1	18.7	22.0	34.4	86	35.9	15.6	18.8	28.7	163
C22	Liver	8	3.4	2.4	2.5	3.2	4	1.7	0.9	1.0	1.4	12
C23, C24	Gallbladder	10	4.3	2.4	2.9	4.5	7	2.9	0.9	1.3	2.2	17
C25	Pancreas	33	14.2	8.6	9.8	13.7	25	10.4	4.3	5.2	8.2	58
C30, C31	Nasal cavities	3	1.3	0.9	1.0	1.1	0	0.0	0.0	0.0	0.0	3
C32	Larynx	2	0.9	0.5	0.6	0.9	0	0.0	0.0	0.0	0.0	2
C33, C34	Lung	152	65.3	36.7	43.2	64.6	112	46.7	25.6	28.9	40.0	264
C37, C38	Thymus etc	1	0.4	0.3	0.3	0.4	0	0.0	0.0	0.0	0.0	1
C40, C41	Bone	1	0.4	0.3	0.4	0.4	0	0.0	0.0	0.0	0.0	1
C43	Melanoma of skin	24	10.3	5.9	7.1	10.3	5	2.1	0.5	0.7	1.4	29
C44	Skin	3	1.3	0.8	0.9	1.2	1	0.4	0.1	0.2	0.3	4
C45	Mesothelioma	14	6.0	3.6	4.1	5.6	3	1.3	0.8	0.8	1.1	17
C47, C49	Connective tissue	7	3.0	3.0	3.0	3.0	4	1.7	1.1	1.2	1.4	11
C50	Breast	1	0.4	0.3	0.3	0.4	69	28.8	17.7	19.6	25.3	70
C53	Cervix	-	-	-	-	-	15	6.3	4.0	4.4	5.6	15
C54, C55	Uterus	-	-	-	-	-	7	2.9	1.9	2.0	2.5	7
C51, C52, C57	Vulva etc	-	-	-	-	-	2	0.8	0.4	0.5	0.6	2
C56	Ovary	-	-	-	-	-	22	9.2	5.1	5.8	7.8	22
C58	Placenta	-	-	-	-	-	0	0	0	0	0	0
C61	Prostate	95	40.8	21.5	26.7	44.8	-	-	-	-	-	95
C62	Testis	0	0	0	0	0	-	-	-	-	-	0
C60, C63	Penis etc	0	0	0	0	0	-	-	-	-	-	0
C64	Kidney	14	6.0	3.4	4.1	6.0	6	2.5	1.2	1.4	2.1	20
C67	Bladder	18	7.7	3.8	4.9	8.0	4	1.7	0.4	0.7	1.3	22
C65, C66, C68	Renal pelvis etc	2	0.9	0.5	0.6	0.9	4	1.7	0.7	0.8	1.3	6
C69	Eye	1	0.4	0.4	0.5	0.5	0	0.0	0.0	0.0	0.0	1
C71	Brain	16	6.9	4.4	4.9	6.5	8	3.3	2.1	2.2	2.9	24
C70, C72	Central nervous system	0	0	0	0	0	0	0	0	0	0	0
C73	Thyroid	1	0.4	0.3	0.3	0.4	5	2.1	1.2	1.3	1.8	6
C74, C75	Other endocrine	1	0.4	0.4	0.5	0.5	0	0.0	0.0	0.0	0.0	1
C81 - C85	All lymphomas	24	10.3	6.3	7.1	10.2	19	7.9	3.3	4.0	6.3	43
C90	Multiple myeloma	4	1.7	1.0	1.2	1.8	7	2.9	1.1	1.4	2.2	11
C91 - C95	All leukaemia	24	10.3	7.3	8.0	10.4	17	7.1	2.2	3.0	5.3	41
C80	Unspecified sites	37	15.9	8.7	10.5	16.4	27	11.3	4.2	5.3	8.6	64
<b>C00 - C96</b>	<b>All cancers (excl. NMSC)</b>	<b>639</b>	<b>274.3</b>	<b>158.7</b>	<b>186.3</b>	<b>279.1</b>	<b>489</b>	<b>204.0</b>	<b>102.0</b>	<b>118.1</b>	<b>169.1</b>	<b>1128</b>

